# New York State Expanded Behavioral Risk Factor Surveillance System

**Nassau County** 

Final Report July 2008 - June 2009

New York State Department of Health December 2009

Introduction	12
Table 1. Nassau County and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in Nassau County and New York State	14
Table 2b. Fair or poor general health among adults in Nassau County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Nassau County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Nassau County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Nassau County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Nassau County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Nassau County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Nassau County and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in Nassau County and New York State	22
Table 6b. Health insurance among adults in Nassau County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Nassau County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Nassau County and New York State - Age-adjusted	25

Table 8a. No health ins	surance among adults in Nassau County and New York State	26
Table 8b. No health ins State - Age-a	surance among adults in Nassau County and New York adjusted	27
Table 9a. No health ins	surance among adults aged 18-64 in Nassau County and New York State	28
Table 9b. No health ins State - Age-a	surance among adults aged 18-64 in Nassau County and New York adjusted	29
Table 10a. Adults with	regular health care provider in Nassau County and New York State	30
Table 10b. Adults with State - Age	regular health care provider in Nassau County and New York -adjusted	31
Table 11a. Cost preven	ated visit to doctor within the past year among adults in Nassau County ork State	32
	nted visit to doctor within the past year among adults in Nassau County ork State - Age-adjusted	33
	octor for routine checkup within the past year among adults in Nassau New York State	34
	octor for routine checkup within the past year among adults in Nassau New York State - Age-adjusted	35
	octor for routine checkup within the past 2 years among adults in Nassau New York State	36
	octor for routine checkup within the past 2 years among adults in Nassau New York State - Age-adjusted	37
Table 14a. Dentist visi	t within the past year among adults in Nassau County and New York State	38
Table 14b. Dentist visi State - Age	t within the past year among adults in Nassau County and New York -adjusted	39
Table 15a. Had teeth c	leaned within the past year among adults in Nassau County and New York	40
Table 15b. Had teeth c State - Age	leaned within the past year among adults in Nassau County and New York -adjusted	41

Table 16a	Permanent teeth extracted due to decay or gum disease among adults in Nassau County and New York State	42
Table 16b	Permanent teeth extracted due to decay or gum disease among adults in Nassau County and New York State - Age-adjusted	43
Table 17a	Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Nassau County and New York State	44
Table 17b	. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Nassau County and New York State - Age-adjusted	45
Table 18a	Flu shot within the past 12 months in Nassau County and New York State	46
Table 18b	Flu shot within the past 12 months in Nassau County and New York State - Age-adjusted	47
Table 19a	Flu vaccine that was sprayed in nose within the past 12 months in Nassau County and New York State	48
Table 19b	. Flu vaccine that was sprayed in nose within the past 12 months in Nassau County and New York State - Age-adjusted	49
Table 20a	Flu shot or flu vaccine sprayed in nose within the past 12 months in Nassau County and New York State	50
Table 20b	Flu shot or flu vaccine sprayed in nose within the past 12 months in Nassau County and New York State - Age-adjusted	51
Table 21a	Flu shot among adults aged 65 and older in Nassau County and New York State	52
Table 21b	Flu shot among adults aged 65 and older in Nassau County and New York State - Age-adjusted	53
Table 22a	Pneumonia shot or pneumococcal vaccine in Nassau County and New York State	54
Table 22b	Pneumonia shot or pneumococcal vaccine in Nassau County and New York State - Age-adjusted	55
Table 23a.	Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Nassau County and New York State	56
Table 23b	Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Nassau County and New York State - Age-adjusted	57

# **Cancer Screening**

Table 24a.	Ever had a mammogram among women aged 40 and older in Nassau County and New York State	58
Table 24b	. Ever had a mammogram among women aged 40 and older in Nassau County and New York State - Age-adjusted	59
Table 25a.	Had mammogram within the past 2 years among women aged 40 and older in Nassau County and New York State	60
Table 25b	. Had mammogram within the past 2 years among women aged 40 and older in Nassau County and New York State - Age-adjusted	61
Table 26a.	Had mammogram within the past 2 years among women aged 50 and older in Nassau County and New York State	62
Table 26b	. Had mammogram within the past 2 years among women aged 50 and older in Nassau County and New York State - Age-adjusted	63
Table 27a.	Ever had a Pap test among women in Nassau County and New York State	64
Table 27b	Ever had a Pap test among women in Nassau County and New York State - Age-adjusted	65
Table 28a.	. Had a Pap test within the past 3 years among women in Nassau County and New York State	66
Table 28b	. Had a Pap test within the past 3 years among women in Nassau County and New York State - Age-adjusted	67
Table 29a.	Ever had a digital rectal exam among men aged 40 and older in Nassau County and New York State	68
Table 29b	. Ever had a digital rectal exam among men aged 40 and older in Nassau County and New York State - Age-adjusted	69
Table 30a.	Had digital rectal exam within the past 2 years among men aged 40 and older in Nassau County and New York State	70
Table 30b	. Had digital rectal exam within the past 2 years among men aged 40 and older in Nassau County and New York State - Age-adjusted	71

Table 31a.	Ever had a prostate specific antigen test among men aged 40 and older in Nassau County and New York State	72
Table 31b.	Ever had a prostate specific antigen test among men aged 40 and older in Nassau County and New York State - Age-adjusted	73
Table 32a.	Had prostate specific antigen test within the past 2 years among men aged 40 and older in Nassau County and New York State	74
Table 32b.	Had prostate specific antigen test within the past 2 years among men aged 40 and older in Nassau County and New York State - Age-adjusted	75
Table 33a.	Prostate cancer among men aged 40 and older in Nassau County and New York State	76
Table 33b.	Prostate cancer among men aged 40 and older in Nassau County and New York State - Age-adjusted	77
Table 34a.	Home blood stool test ever used among adults aged 50 and older in Nassau County and New York State	78
Table 34b.	Home blood stool test ever used among adults aged 50 and older in Nassau County and New York State - Age-adjusted	79
Table 35a.	Home blood stool test used within the past year among adults aged 50 and older in Nassau County and New York State	80
Table 35b.	Home blood stool test used within the past year among adults aged 50 and older in Nassau County and New York State - Age-adjusted	81
Table 36a.	Home blood stool test used within the past 2 years among adults aged 50 and older in Nassau County and New York State	82
Table 36b.	Home blood stool test used within the past 2 years among adults aged 50 and older in Nassau County and New York State - Age-adjusted	83
Table 37a.	Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Nassau County and New York State	84
Table 37b.	Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Nassau County and New York State - Age-adjusted	85
Table 38a.	Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Nassau County and New York State	86

Table 38b	. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Nassau County and New York State - Age-adjusted	87
Table 39a.	Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Nassau County and New York State	88
Table 39b	Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Nassau County and New York State - Age-adjusted	89
Chronic	Diseases	
Table 40a.	High blood pressure among adults in Nassau County and New York State	90
Table 40b	. High blood pressure among adults in Nassau County and New York State - Age-adjusted	91
Table 41a.	Blood pressure medication among adults with high blood pressure in Nassau County and New York State	92
Table 41b	Blood pressure medication among adults with high blood pressure in Nassau County and New York State - Age-adjusted	93
Table 42a.	Ever had blood cholesterol checked among adults in Nassau County and New York State	94
Table 42b	Ever had blood cholesterol checked among adults in Nassau County and New York State - Age-adjusted	95
Table 43a.	Blood cholesterol checked within the past 5 years among adults in Nassau County and New York State	96
Table 43b	Blood cholesterol checked within the past 5 years among adults in Nassau County and New York State - Age-adjusted	97
Table 44a.	Cardiovascular disease among adults in Nassau County and New York State	98
Table 44b	. Cardiovascular disease among adults in Nassau County and New York State - Age-adjusted	99
Table 45a.	Coronary heart disease among adults in Nassau County and New York State	100

Table 45b.	Coronary heart disease among adults in Nassau County and New York State - Age-adjusted	101
Table 46a.	Ever diagnosed with diabetes among adults in Nassau County and New York State	102
Table 46b.	Ever diagnosed with diabetes among adults in Nassau County and New York State - Age-adjusted	103
Table 47a.	Ever diagnosed with asthma (lifetime) among adults in Nassau County and New York State	104
Table 47b.	Ever diagnosed with asthma (lifetime) among adults in Nassau County and New York State - Age-adjusted	105
Table 48a.	Current asthma among adults in Nassau County and New York State	106
Table 48b.	Current asthma among adults in Nassau County and New York State - Age-adjusted	107
Table 49a.	Chronic joint symptoms among adults in Nassau County and New York State	108
Table 49b.	Chronic joint symptoms among adults in Nassau County and New York State - Age-adjusted	109
Table 50a.	Arthritis among adults in Nassau County and New York State	110
Table 50b.	Arthritis among adults in Nassau County and New York State - Age-adjusted	111
Table 51a.	Activity limitations because of physical, mental or emotional problems among adults in Nassau County and New York State	112
Table 51b.	Activity limitations because of physical, mental or emotional problems among adults in Nassau County and New York State - Age-adjusted	113
Table 52a.	Have health problem that requires the use of special equipment among adults in Nassau County and New York State	114
Table 52b.	Have health problem that requires the use of special equipment among adults in Nassau County and New York State - Age-adjusted	115
Table 53a.	Disability among adults in Nassau County and New York State	116
Table 53b.	Disability among adults in Nassau County and New York State - Age-adjusted	117

#### **Health Behaviors**

Table 54a.	Current smoking among adults in Nassau County and New York State	118
Table 54b.	Current smoking among adults in Nassau County and New York State - Age-adjusted	119
Table 55a.	Everyday smoking among adults in Nassau County and New York State	120
Table 55b.	Everyday smoking among adults in Nassau County and New York State - Age-adjusted	121
Table 56a.	Adults living in homes in which smoking is prohibited in Nassau County and New York State	122
Table 56b.	Adults living in homes in which smoking is prohibited in Nassau County and New York State - Age-adjusted	123
Table 57a.	Binge drinking within the past month among adults in Nassau County and New York State	124
Table 57b.	Binge drinking within the past month among adults in Nassau County and New York State - Age-adjusted	125
Table 58a.	Heavy drinking within the past month among adults in Nassau County and New York State	126
Table 58b.	Heavy drinking within the past month among adults in Nassau County and New York State - Age-adjusted	127
Table 59a.	Never or rarely been asked about sexual history during routine check-up among adults in Nassau County and New York State	128
Table 59b.	Never or rarely been asked about sexual history during routine check-up among adults in Nassau County and New York State - Age-adjusted	129
Table 60a.	Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Nassau County and New York State	130
Table 60b.	Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Nassau County and New York State - Age-adjusted	131

Table 61a.	Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Nassau County and New York State	132
Table 61b.	Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Nassau County and New York State - Age-adjusted	133
Overwei	ght / Obesity	
Table 62a.	Overweight among adults in Nassau County and New York State	134
Table 62b.	Overweight among adults in Nassau County and New York State - Age-adjusted	135
Table 63a.	Obesity among adults in Nassau County and New York State	136
Table 63b.	Obesity among adults in Nassau County and New York State - Age-adjusted	137
Table 64a.	Overweight or obesity among adults in Nassau County and New York State	138
Table 64b.	Overweight or obesity among adults in Nassau County and New York State - Age-adjusted	139
Table 65a.	Received advice about weight by a health professional among adults in Nassau County and New York State	140
Table 65b.	Received advice about weight by a health professional among adults in Nassau County and New York State - Age-adjusted	141
Table 66a.	Received advice to lose weight by a health professional among adults provided weight advice in Nassau County and New York State	142
Table 66b.	Received advice to lose weight by a health professional among adults provided weight advice in Nassau County and New York State - Age-adjusted	143
Physical	Activity / Nutrition	
Table 67a.	No leisure-time physical activity among adults in Nassau County and New York State	144
Table 67b.	No leisure-time physical activity among adults in Nassau County and New York State - Age-adjusted	145

Table 68a.	Consumption of 5 or more servings of fruits and vegetables per day among adults in Nassau County and New York State	146
Table 68b	Consumption of 5 or more servings of fruits and vegetables per day among adults in Nassau County and New York State - Age-adjusted	147
Injury		
Table 69a.	Had a fall within the past 3 months among adults in Nassau County and New York State	148
Table 69b	. Had a fall within the past 3 months among adults in Nassau County and New York State - Age-adjusted	149
Table 70a.	Had a fall that caused injury within the past 3 months among adults in Nassau County and New York State	150
Table 70b	. Had a fall that caused injury within the past 3 months among adults in Nassau County and New York State - Age-adjusted	151

#### Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

Table 1. Nassau County and New York State adult US Census Bureau population estimates,<sup>1</sup> 2007

	Nassau C	ounty	New York State			
	Adult Population <sup>1</sup>	Percent	Adult Population <sup>1</sup>	Percent		
Total	1,005,031	100	14,884,315	100		
Gender						
Male	480,140	47.8	7,103,791	47.7		
Female	524,891	52.2	7,780,524	52.3		
Age (years)						
18-34	243,193	24.2	4,497,479	30.2		
35-44	181,849	18.1	2,844,115	19.1		
45-54	217,022	21.6	2,854,316	19.2		
55-64	168,348	16.8	2,142,000	14.4		
≥ 65	194,619	19.4	2,546,405	17.1		
Race/Ethnicity						
White Non-Hispanic	713,320	71.0	9,343,884	62.8		
Black Non-Hispanic	104,418	10.4	2,156,941	14.5		
Hispanic	115,209	11.5	2,258,717	15.2		
Other Non-Hispanic	72,084	7.2	1,124,773	7.6		

<sup>&</sup>lt;sup>1</sup> Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged\_race.htm as of November 25, 2009.

Table 2a. Fair or poor general health among adults in Nassau County and New York State

	Nassau County				New York State			
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	647	153,600	15.3	(11.5 - 19.2)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	243	60,500	12.7	(6.7 - 18.7)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	404	93,100	17.8	(12.8 - 22.7)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	*	*	*	*	4,985	289,200	10.0	(6.8 - 13.3)
45-54	115	31,800	14.7	(7.0 - 22.3)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	137	29,400	17.5	(7.8 - 27.2)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	220	49,000	25.2	(18.4 - 31.9)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	554	85,800	12.1	(9.0 - 15.1)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	156	70,300	26.5	(16.7 - 36.3)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	145	31,700	13.4	(6.0 - 20.7)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	343	51,600	10.4	(6.1 - 14.7)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	*	*	*	*	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	*	*	*	*	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	109	26,900	16.7	(7.4 - 26.1)	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	275	32,100	7.3	(3.8 - 10.8)	8,562	218,300	4.9	(3.6 - 6.1)

 $<sup>^{1}</sup>$  Defined as reporting fair or poor general health

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a

Table 2b. Fair or poor general health among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	1	Nassau C	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	647	13.8	(9.9 - 17.6)	38,124	16.7	(14.8 - 18.6)		
Gender								
Male	243	12.8	(6.4 - 19.3)	14,599	15.3	(12.6 - 18.0)		
Female	404	14.7	(10.2 - 19.1)	23,525	18.0	(15.2 - 20.7)		
Race/Ethnicity								
White Non-Hispanic	554	10.8	(7.9 - 13.7)	35,811	12.9	(11.4 - 14.4)		
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)		
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)		
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)		
Education								
≤ High School	*	*	*	15,682	25.8	(22.1 - 29.5)		
Some College	145	11.7	(5.0 - 18.3)	9,904	14.1	(10.7 - 17.6)		
≥ College Degree	343	9.3	(5.0 - 13.5)	12,457	8.5	(6.3 - 10.8)		
Income								
≤ \$24,999	*	*	*	9,633	35.9	(29.7 - 42.0)		
\$25,000-\$49,999	*	*	*	9,649	15.0	(11.6 - 18.5)		
\$50,000-\$74,999	109	13.3	(6.0 - 20.6)	5,769	11.8	(7.7 - 16.0)		
≥ \$75,000	275	6.8	(3.4 - 10.1)	8,562	4.7	(3.5 - 6.0)		

<sup>&</sup>lt;sup>1</sup> Defined as reporting fair or poor general health

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 3a. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	y		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	635	86,200	8.7	(6.2 - 11.2)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	239	38,100	8.1	(4.2 - 12.0)	14,325	604,200	8.7	(7.0 - 10.5)
Female	396	48,100	9.3	(6.1 - 12.5)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	*	*	*	*	4,948	232,100	8.1	(5.2 - 11.1)
45-54	113	19,000	9.0	(3.5 - 14.4)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	134	18,200	11.1	(3.4 - 18.9)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	212	32,800	17.4	(11.4 - 23.5)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	543	65,200	9.3	(6.7 - 11.9)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	149	21,900	8.6	(3.6 - 13.5)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	145	25,200	10.7	(4.6 - 16.7)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	338	39,100	7.9	(4.7 - 11.1)	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	68	10,400	8.8	(1.6 - 15.9)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	86	15,800	12.8	(5.4 - 20.2)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	106	22,500	14.2	(5.0 - 23.5)	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	274	27,200	6.2	(3.2 - 9.1)	8,510	219,100	4.9	(3.6 - 6.2)

 $<sup>^{\</sup>mathrm{1}}$  Defined as 14 or more days within the past month when physical health was not good

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 3b. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	635	7.7	(5.5 - 10.0)	37,327	9.9	(8.4 - 11.4)		
Gender								
Male	239	7.9	(4.1 - 11.7)	14,325	8.7	(6.9 - 10.5)		
Female	396	7.6	(4.9 - 10.2)	23,002	11.0	(8.7 - 13.2)		
Race/Ethnicity								
White Non-Hispanic	543	8.7	(6.0 - 11.3)	35,073	9.9	(8.5 - 11.3)		
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)		
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)		
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)		
Education								
≤ High School	149	7.7	(3.0 - 12.3)	15,178	12.7	(10.2 - 15.1)		
Some College	145	8.7	(4.0 - 13.5)	9,751	10.1	(7.2 - 12.9)		
≥ College Degree	338	7.3	(4.2 - 10.5)	12,327	6.3	(4.7 - 8.0)		
Income								
≤ \$24,999	68	8.4	(0.9 - 15.8)	9,294	18.4	(13.5 - 23.4)		
\$25,000-\$49,999	86	10.6	(4.1 - 17.1)	9,495	9.4	(7.0 - 11.9)		
\$50,000-\$74,999	106	11.2	(4.2 - 18.2)	5,733	9.0	(5.3 - 12.6)		
≥ \$75,000	274	5.9	(2.9 - 9.0)	8,510	4.9	(3.5 - 6.2)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as 14 or more days within the past month when physical health was not good

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 4a. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au County	/		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	94,600	9.5	(6.4 - 12.6)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	240	30,600	6.5	(2.4 - 10.6)	14,407	555,200	7.9	(6.3 - 9.6)
Female	403	64,000	12.2	(7.6 - 16.9)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	*	*	*	*	4,947	254,100	8.9	(6.7 - 11.1)
45-54	115	34,000	15.7	(6.7 - 24.6)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	135	14,000	8.5	(3.5 - 13.4)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	217	11,100	5.8	(2.4 - 9.3)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	552	67,400	9.5	(6.3 - 12.8)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	155	31,000	11.7	(5.6 - 17.9)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	144	27,800	12.0	(3.1 - 20.8)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	341	35,900	7.2	(3.9 - 10.5)	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	*	*	*	*	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	*	*	*	*	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	107	19,900	12.5	(4.4 - 20.7)	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	275	39,100	8.9	(4.3 - 13.4)	8,520	276,500	6.2	(4.6 - 7.7)

 $<sup>^{\</sup>mathrm{1}}$  Defined as 14 or more days within the past month when mental health was not good

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 4b. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau Co	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	9.6	(6.2 - 13.1)	37,528	10.2	(8.7 - 11.7)
Gender						
Male	240	5.8	(2.2 - 9.3)	14,407	7.9	(6.3 - 9.6)
Female	403	13.1	(7.4 - 18.8)	23,121	12.2	(9.8 - 14.6)
Race/Ethnicity						
White Non-Hispanic	552	10.0	(6.1 - 13.9)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
Education						
≤ High School	155	10.3	(4.8 - 15.9)	15,335	12.3	(10.0 - 14.6)
Some College	*	*	*	9,771	11.8	(8.5 - 15.1)
≥ College Degree	341	7.7	(3.7 - 11.6)	12,342	6.3	(4.6 - 8.0)
Income						
≤ \$24 <b>,</b> 999	*	*	*	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	*	*	*	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	107	13.7	(4.2 - 23.2)	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	275	8.2	(3.9 - 12.5)	8,520	6.2	(4.6 - 7.7)

 $<sup>^{\</sup>rm 1}$  Defined as 14 or more days within the past month when mental health was not good

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 5a. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	633	166,100	16.9	(13.1 - 20.6)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	237	63,600	13.6	(8.2 - 19.0)	14,231	937,700	13.6	(11.4 - 15.8)
Female	396	102,500	19.8	(14.6 - 25.0)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	108	18,600	10.2	(4.1 - 16.4)	4,924	388,300	13.7	(10.4 - 17.0)
45-54	115	47,100	21.7	(12.4 - 31.1)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	133	27,800	17.2	(8.7 - 25.6)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	210	40,800	22.0	(15.4 - 28.5)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	542	124,600	17.8	(13.9 - 21.8)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	150	46,800	18.1	(10.8 - 25.3)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	143	51,300	22.2	(12.3 - 32.1)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	337	68,000	13.8	(9.5 - 18.1)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	*	*	*	*	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	*	*	*	*	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	*	*	*	*	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	274	61,500	14.0	(8.8 - 19.1)	8,475	447,600	10.0	(8.1 - 12.0)

 $<sup>^{1}</sup>$  Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation <sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 5b. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau C	ounty	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	633	16.3	(12.3 - 20.3)	37,005	16.4	(14.7 - 18.2)		
Gender								
Male	237	12.8	(7.8 - 17.8)	14,231	13.6	(11.4 - 15.8)		
Female	396	19.4	(13.4 - 25.4)	22,774	19.0	(16.3 - 21.6)		
Race/Ethnicity								
White Non-Hispanic	542	17.8	(13.4 - 22.3)	34,778	16.3	(14.7 - 17.8)		
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)		
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)		
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)		
Education								
≤ High School	150	16.4	(9.4 - 23.3)	15,018	20.1	(17.1 - 23.1)		
Some College	*	*	*	9,672	17.6	(14.1 - 21.1)		
≥ College Degree	337	13.8	(9.0 - 18.7)	12,244	11.4	(9.2 - 13.7)		
Income								
≤ \$24,999	*	*	*	9,199	27.6	(22.1 - 33.1)		
\$25,000-\$49,999	*	*	*	9,425	16.1	(12.9 - 19.3)		
\$50,000-\$74,999	*	*	*	5,696	15.6	(11.0 - 20.1)		
≥ \$75,000	274	13.2	(8.1 - 18.3)	8,475	10.0	(8.0 - 12.0)		

 $<sup>^{1}</sup>$  Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 6a. Health insurance among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	647	912,200	90.9	(87.2 - 94.6)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	244	418,500	87.2	(80.7 - 93.7)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	403	493,600	94.4	(90.8 - 98.0)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	109	166,000	91.3	(84.4 - 98.2)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	114	204,900	95.0	(90.0 - 100)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	137	152,900	90.8	(83.4 - 98.3)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	219	191,000	98.4	(95.3 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	554	681,700	95.8	(93.4 - 98.2)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	156	223,100	84.2	(76.1 - 92.2)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	145	211,700	89.6	(81.5 - 97.6)	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	343	474,700	95.1	(90.4 - 99.8)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	*	*	*	*	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	109	148,100	92.3	(84.0 - 100)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	276	439,400	99.1	(98.0 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare <sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have health insurance (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 6b. Health insurance among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	647	90.6	(86.6 - 94.6)	38,119	86.7	(84.4 - 89.1)		
Gender								
Male	244	87.3	(80.5 - 94.1)	14,597	83.6	(79.7 - 87.5)		
Female	403	93.6	(89.2 - 98.1)	23,522	89.6	(86.8 - 92.3)		
Race/Ethnicity								
White Non-Hispanic	554	95.4	(92.4 - 98.4)	35,811	92.1	(90.5 - 93.7)		
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)		
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	156	84.4	(76.0 - 92.8)	15,673	80.1	(75.4 - 84.7)		
Some College	145	88.5	(79.3 - 97.7)	9,906	84.6	(79.7 - 89.4)		
≥ College Degree	343	94.7	(89.4 - 100)	12,459	95.4	(93.6 - 97.2)		
Income								
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)		
\$25,000-\$49,999	*	*	*	9,662	81.9	(76.0 - 87.9)		
\$50,000-\$74,999	109	92.8	(85.0 - 100)	5,772	94.0	(91.3 - 96.7)		
≥ \$75,000	276	99.3	(98.4 - 100)	8,567	97.5	(96.6 - 98.5)		

 $<sup>^1</sup>$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 7a. Health insurance among adults aged 18-64<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	423	717,400	89.1	(84.6 - 93.6)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	164	339,600	85.3	(77.7 - 93.0)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	259	377,700	92.8	(88.2 - 97.4)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	109	166,000	91.3	(84.4 - 98.2)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	114	204,900	95.0	(90.0 - 100)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	132	149,100	90.6	(83.0 - 98.2)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	342	501,700	94.4	(91.2 - 97.5)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	99	169,500	87.3	(77.5 - 97.0)	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	239	383,600	94.0	(88.3 - 99.7)	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	*	*	*	*	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	*	*	*	*	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	223	389,700	99.0	(97.8 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

 $<sup>^{1}</sup>$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 7b. Health insurance among adults aged 18-64<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	1	lassau C	ounty	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	423	89.0	(84.2 - 93.8)	25,165	84.3	(81.5 - 87.2)		
Gender								
Male	164	85.4	(77.4 - 93.5)	10,078	80.5	(75.9 - 85.1)		
Female	259	92.3	(87.0 - 97.6)	15,087	87.9	(84.6 - 91.2)		
Race/Ethnicity								
White Non-Hispanic	342	94.1	(90.3 - 97.9)	23,354	90.3	(88.3 - 92.3)		
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	9,037	75.1	(69.3 - 80.9)		
Some College	*	*	*	7,031	82.6	(77.1 - 88.1)		
≥ College Degree	239	93.7	(87.5 - 99.9)	9,068	94.6	(92.5 - 96.8)		
Income								
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)		
\$25,000-\$49,999	*	*	*	6,180	77.5	(70.2 - 84.7)		
\$50,000-\$74,999	70	90.7	(80.7 - 100)	4,554	93.0	(89.8 - 96.2)		
≥ \$75,000	223	99.2	(98.2 - 100)	7,324	97.3	(96.3 - 98.3)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 8a. No health insurance among adults in Nassau County and New York State

		Nassa	au County	<i>'</i>		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	647	91,100	9.1	(5.4 - 12.8)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	244	61,600	12.8	(6.3 - 19.3)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	403	29,500	5.6	(2.0 - 9.2)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	*	*	*	*	7,831	316,500	11.1	(6.8 - 15.4)
55-64	*	*	*	*	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	554	29,800	4.2	(1.8 - 6.6)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	156	41,900	15.8	(7.8 - 23.9)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	*	*	*	*	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

<sup>&</sup>lt;sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare <sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have no health insurance (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 8b. No health insurance among adults 1 in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	647	9.4	(5.4 - 13.4)	38,119	13.3	(10.9 - 15.6)		
Gender								
Male	244	12.7	(5.9 - 19.5)	14,597	16.4	(12.5 - 20.3)		
Female	403	6.4	(1.9 - 10.8)	23,522	10.4	(7.7 - 13.2)		
Race/Ethnicity								
White Non-Hispanic	554	4.6	(1.6 - 7.6)	35,811	7.9	(6.3 - 9.5)		
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)		
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	156	15.6	(7.2 - 24.0)	15,673	19.9	(15.3 - 24.6)		
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)		
≥ College Degree	*	*	*	12,459	4.6	(2.8 - 6.4)		
Income								
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)		
\$25,000-\$49,999	*	*	*	9,662	18.1	(12.1 - 24.0)		
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)		
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)		

<sup>&</sup>lt;sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 9a. No health insurance among adults aged 18-641 in Nassau County and New York State

		Nassa	au County	/		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	423	88,000	10.9	(6.4 - 15.4)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	164	58,500	14.7	(7.0 - 22.3)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	259	29,500	7.2	(2.6 - 11.8)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	*	*	*	*	7,791	316,100	11.3	(7.0 - 15.7)
55-64	*	*	*	*	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	342	29,800	5.6	(2.5 - 8.8)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

<sup>&</sup>lt;sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64 <sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 9b. No health insurance among adults aged 18-64<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	423	11.0	(6.2 - 15.8)	25,165	15.7	(12.8 - 18.5)		
Gender								
Male	164	14.6	(6.5 - 22.6)	10,078	19.5	(14.9 - 24.1)		
Female	259	7.7	(2.4 - 13.0)	15,087	12.1	(8.8 - 15.4)		
Race/Ethnicity								
White Non-Hispanic	342	5.9	(2.1 - 9.7)	23,354	9.7	(7.7 - 11.7)		
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	9,037	24.9	(19.1 - 30.7)		
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)		
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)		
Income								
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)		
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)		
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)		
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)		

<sup>&</sup>lt;sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64 <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 10a. Adults with regular health care provider in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	648	884,400	88.1	(84.0 - 92.1)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	243	414,600	86.5	(80.4 - 92.5)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	405	469,800	89.5	(83.9 - 95.1)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	109	166,300	91.4	(85.0 - 97.9)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	115	198,100	91.3	(85.3 - 97.3)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	137	154,900	92.0	(86.9 - 97.0)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	219	188,500	97.2	(94.6 - 99.7)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	555	651,900	91.5	(88.1 - 94.8)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	146	211,200	88.9	(81.4 - 96.3)	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	344	447,600	89.6	(84.9 - 94.3)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	*	*	*	*	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	109	153,700	95.8	(91.5 - 100)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	276	396,200	89.4	(84.3 - 94.4)	8,560	4,122,800	91.9	(90.2 - 93.7)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

nearest hundred)
<sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 10b. Adults with regular health care provider<sup>1</sup> in Nassau County and New York
State - Age-Adjusted<sup>2</sup>

	N	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	648	86.9	(82.1 - 91.7)	38,135	83.0	(80.4 - 85.5)		
Gender								
Male	243	85.9	(79.4 - 92.4)	14,593	78.1	(73.9 - 82.3)		
Female	405	87.8	(80.9 - 94.8)	23,542	87.4	(84.6 - 90.1)		
Race/Ethnicity								
White Non-Hispanic	555	90.2	(85.9 - 94.5)	35,821	88.0	(86.2 - 89.7)		
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)		
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,692	75.8	(71.0 - 80.6)		
Some College	146	86.6	(77.2 - 95.9)	9,904	82.9	(77.5 - 88.3)		
≥ College Degree	344	88.8	(83.2 - 94.3)	12,456	89.8	(87.1 - 92.4)		
Income								
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)		
\$25,000-\$49,999	*	*	*	9,655	77.4	(70.5 - 84.3)		
\$50,000-\$74,999	109	95.1	(89.8 - 100)	5,771	89.6	(85.3 - 94.0)		
≥ \$75,000	276	88.4	(82.5 - 94.3)	8,560	91.6	(89.8 - 93.4)		

 $<sup>^{1}</sup>$  Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 11a. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au County	,		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	646	122,000	12.2	(8.1 - 16.2)	38,143	2,034,800	13.7	(11.6 - 15.8)		
Gender										
Male	244	43,500	9.1	(4.0 - 14.1)	14,607	903,300	12.7	(9.5 - 16.0)		
Female	402	78,500	15.0	(8.9 - 21.1)	23,536	1,131,500	14.6	(11.8 - 17.4)		
Age (years)										
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)		
35-44	107	21,400	11.9	(4.2 - 19.6)	4,987	347,900	12.1	(8.0 - 16.2)		
45-54	114	21,500	9.9	(3.5 - 16.4)	7,825	382,200	13.5	(9.9 - 17.1)		
55-64	137	15,200	9.0	(3.1 - 14.9)	8,766	236,800	11.1	(8.0 - 14.2)		
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)		
Race/Ethnicity										
White Non-Hispanic	553	43,900	6.2	(3.4 - 8.9)	35,833	984,600	10.6	(8.9 - 12.2)		
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)		
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)		
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)		
Education										
≤ High School	*	*	*	*	15,696	996,600	17.8	(13.8 - 21.9)		
Some College	146	29,500	12.4	(4.6 - 20.3)	9,911	512,400	14.5	(10.4 - 18.6)		
≥ College Degree	342	45,300	9.1	(4.8 - 13.4)	12,453	466,900	8.3	(6.0 - 10.6)		
Income										
≤ \$24,999	*	*	*	*	9,639	854,500	24.3	(18.4 - 30.3)		
\$25,000-\$49,999	*	*	*	*	9,656	532,900	18.4	(12.5 - 24.3)		
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)		
≥ \$75 <b>,</b> 000	273	14,400	3.3	(1.2 - 5.3)	8,561	192,000	4.3	(2.9 - 5.6)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as needing to see a doctor within the past year but not being able to because of cost

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 11b. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	646	12.9	(8.3 - 17.5)	38,143	13.8	(11.7 - 15.9)		
Gender								
Male	244	9.4	(4.0 - 14.8)	14,607	12.5	(9.4 - 15.6)		
Female	402	16.1	(9.0 - 23.3)	23,536	15.0	(12.1 - 17.9)		
Race/Ethnicity								
White Non-Hispanic	553	7.3	(3.7 - 10.9)	35,833	10.8	(9.2 - 12.5)		
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)		
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)		
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)		
Education								
≤ High School	*	*	*	15,696	17.8	(13.8 - 21.7)		
Some College	146	14.8	(5.3 - 24.2)	9,911	15.3	(11.0 - 19.5)		
≥ College Degree	342	8.8	(4.2 - 13.5)	12,453	8.2	(6.0 - 10.5)		
Income								
≤ \$24,999	*	*	*	9,639	24.5	(18.6 - 30.4)		
\$25,000-\$49,999	*	*	*	9,656	18.7	(12.9 - 24.6)		
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)		
≥ \$75,000	273	3.3	(1.1 - 5.5)	8,561	4.4	(3.1 - 5.8)		

 $<sup>^{\</sup>rm 1}$  Defined as needing to see a doctor within the past year but not being able to because of cost

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 12a. Visited a doctor for routine checkup within the past year among adults in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	777,400	77.6	(73.1 - 82.1)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	243	366,700	76.7	(69.8 - 83.6)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	403	410,700	78.5	(72.6 - 84.3)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	114	168,700	78.0	(69.4 - 86.5)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	137	132,400	78.6	(70.5 - 86.8)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	220	179,400	92.2	(88.4 - 96.0)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	553	553,800	78.0	(73.4 - 82.6)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	155	206,800	78.3	(69.4 - 87.2)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	342	386,300	77.7	(71.8 - 83.6)	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	109	139,100	86.7	(78.8 - 94.7)	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	273	332,200	75.5	(68.9 - 82.1)	8,547	3,176,500	70.7	(67.3 - 74.2)

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 12b. Visited a doctor for routine checkup within the past year among adults in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	646	76.4	(71.3 - 81.5)	38,026	72.7	(70.3 - 75.0)		
Gender								
Male	243	75.9	(68.6 - 83.3)	14,592	67.1	(63.3 - 70.9)		
Female	403	76.8	(69.7 - 83.9)	23,434	77.7	(75.1 - 80.3)		
Race/Ethnicity								
White Non-Hispanic	553	76.9	(71.5 - 82.2)	35,721	71.9	(70.1 - 73.7)		
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)		
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	155	78.4	(69.0 - 87.9)	15,640	74.4	(70.4 - 78.4)		
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)		
≥ College Degree	342	76.6	(69.9 - 83.4)	12,428	71.7	(68.3 - 75.0)		
Income								
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)		
\$25,000-\$49,999	*	*	*	9,631	71.1	(65.3 - 76.9)		
\$50,000-\$74,999	109	86.6	(78.0 - 95.3)	5,765	74.5	(69.2 - 79.8)		
≥ \$75,000	273	73.8	(66.3 - 81.4)	8,547	70.3	(66.9 - 73.8)		

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	886,600	88.5	(85.0 - 92.1)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	243	414,400	86.7	(81.1 - 92.2)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	403	472,200	90.2	(85.7 - 94.7)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	108	150,500	83.2	(74.3 - 92.1)	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	114	198,200	91.6	(86.0 - 97.2)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	137	154,300	91.7	(86.8 - 96.6)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	220	189,700	97.5	(95.4 - 99.6)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	553	629,800	88.7	(85.1 - 92.3)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	155	226,400	85.8	(78.0 - 93.5)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	146	213,000	89.6	(80.8 - 98.4)	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	342	444,600	89.4	(85.3 - 93.5)	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	*	*	*	*	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	109	153,800	95.8	(91.3 - 100)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	273	380,700	86.5	(81.2 - 91.8)	8,547	3,907,000	87.0	(84.8 - 89.2)

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>
Total	646	87.3	(83.2 - 91.5)	38,026	85.7	(83.7 - 87.6)
Gender						
Male	243	86.2	(80.2 - 92.2)	14,592	81.9	(78.4 - 85.3)
Female	403	88.4	(82.6 - 94.1)	23,434	89.1	(87.4 - 90.9)
Race/Ethnicity						
White Non-Hispanic	553	87.1	(82.6 - 91.6)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	155	86.0	(77.9 - 94.2)	15,640	84.6	(81.1 - 88.1)
Some College	*	*	*	9,880	84.4	(79.8 - 89.0)
≥ College Degree	342	88.0	(83.0 - 93.0)	12,428	87.3	(85.0 - 89.6)
Income						
≤ \$24,999	*	*	*	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	*	*	*	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	109	95.8	(90.8 - 100)	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	273	84.5	(78.3 - 90.8)	8,547	86.4	(84.2 - 88.7)

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 14a. Dentist visit within the past year among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	648	800,700	79.9	(75.5 - 84.2)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	243	359,200	75.2	(67.7 - 82.7)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	405	441,500	84.1	(79.7 - 88.5)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	115	166,800	76.9	(67.8 - 85.9)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	136	147,700	89.1	(83.5 - 94.6)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	220	167,400	86.0	(81.4 - 90.7)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	555	579,600	81.5	(77.5 - 85.5)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	156	198,100	74.8	(65.3 - 84.3)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	145	189,700	80.6	(72.1 - 89.1)	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	344	410,200	82.1	(76.2 - 88.0)	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	*	*	*	*	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	108	138,600	87.8	(81.5 - 94.0)	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	276	377,900	85.2	(80.2 - 90.3)	8,564	3,707,900	82.4	(79.5 - 85.3)

 $<sup>^{\</sup>rm 1}$  Defined as having visited a dentist or a dental clinic for any reason within the past year

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 14b. Dentist visit within the past year among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	648	79.2	(74.4 - 84.0)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	243	73.8	(65.6 - 82.0)	14,573	68.3	(64.6 - 72.1)
Female	405	84.2	(79.3 - 89.0)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	555	81.0	(76.6 - 85.4)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,613	65.3	(60.9 - 69.6)
Some College	145	80.8	(72.0 - 89.5)	9,880	70.4	(65.7 - 75.1)
≥ College Degree	344	80.9	(74.3 - 87.5)	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	108	88.2	(81.6 - 94.8)	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	276	84.3	(78.8 - 89.9)	8,564	82.1	(79.1 - 85.0)

 $<sup>^{\</sup>rm 1}$  Defined as having visited a dentist or a dental clinic for any reason within the past year

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 15a. Had teeth cleaned within the past year among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New York State					
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>			
Total	625	777,500	79.3	(75.0 - 83.6)	34,070	9,985,900	71.7	(69.1 - 74.3)			
Gender											
Male	232	344,700	73.9	(66.4 - 81.4)	13,173	4,616,400	69.0	(65.0 - 73.1)			
Female	393	432,800	84.2	(79.8 - 88.6)	20,897	5,369,500	74.2	(70.8 - 77.5)			
Age (years)											
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)			
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)			
45-54	113	162,800	75.7	(66.4 - 84.9)	7,438	2,022,600	73.3	(69.1 - 77.4)			
55-64	134	131,500	81.1	(71.7 - 90.5)	7,906	1,560,900	77.7	(73.8 - 81.6)			
≥ 65	201	152,800	85.8	(80.6 - 91.0)	9,991	1,543,900	77.1	(73.7 - 80.5)			
Race/Ethnicity											
White Non-Hispanic	534	550,800	79.7	(75.5 - 83.9)	31,982	6,601,500	75.4	(73.4 - 77.4)			
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)			
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)			
Other Non-Hispanic	*	*	*	*	*	*	*	*			
Education											
≤ High School	145	201,500	79.1	(71.0 - 87.2)	12,740	3,388,600	67.7	(63.1 - 72.3)			
Some College	143	177,000	75.6	(66.1 - 85.0)	9,145	2,286,800	68.5	(62.9 - 74.1)			
≥ College Degree	334	397,000	81.2	(75.1 - 87.2)	12,128	4,298,500	78.3	(74.8 - 81.7)			
Income											
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)			
\$25,000-\$49,999	*	*	*	*	8,709	1,681,700	62.6	(55.4 - 69.8)			
\$50,000-\$74,999	102	128,000	84.2	(75.0 - 93.4)	5,534	1,447,800	76.3	(70.7 - 81.8)			
≥ \$75,000	274	360,600	81.8	(76.2 - 87.3)	8,448	3,617,500	80.8	(77.8 - 83.8)			

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 15b. Had teeth cleaned within the past year among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	625	79.3	(74.7 - 83.9)	34,070	71.7	(69.1 - 74.2)
Gender						
Male	232	73.5	(65.5 - 81.4)	13,173	69.2	(65.3 - 73.1)
Female	393	84.6	(80.0 - 89.2)	20,897	73.9	(70.5 - 77.3)
Race/Ethnicity						
White Non-Hispanic	534	79.0	(74.3 - 83.6)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	145	81.0	(73.1 - 89.0)	12,740	67.7	(63.2 - 72.3)
Some College	143	76.9	(67.7 - 86.2)	9,145	68.1	(62.6 - 73.6)
≥ College Degree	334	79.6	(72.8 - 86.4)	12,128	78.3	(74.9 - 81.7)
Income						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	102	86.4	(78.5 - 94.3)	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	274	80.6	(74.4 - 86.7)	8,448	80.4	(77.4 - 83.4)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	640	502,400	50.6	(45.3 - 55.8)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	240	249,100	52.6	(44.4 - 60.8)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	400	253,200	48.8	(42.1 - 55.4)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	*	*	*	*	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	*	*	*	*	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	135	112,300	67.6	(58.2 - 77.1)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	217	143,500	74.6	(68.2 - 81.0)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	549	338,400	48.0	(42.9 - 53.1)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	338	205,500	41.8	(34.8 - 48.8)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	*	*	*	*	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	273	192,900	43.9	(36.2 - 51.6)	8,516	1,738,200	38.8	(35.3 - 42.2)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

 $<sup>^{2}\,\</sup>mathrm{Sample}$  size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred) <sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 16b. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	lassau C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	640	46.4	(40.8 - 52.1)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	240	49.2	(40.5 - 57.8)	14,457	47.9	(44.1 - 51.7)
Female	400	44.0	(36.7 - 51.2)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	549	43.5	(38.0 - 48.9)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	338	38.1	(30.6 - 45.6)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	273	40.8	(32.6 - 49.1)	8,516	37.7	(34.2 - 41.1)

 $<sup>^{\</sup>mathrm{l}}$  Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Nassau County and New York State

		Nassa	au County	•		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	217	15,800	8.2	(4.3 - 12.2)	12,339	491,200	19.6	(16.0 - 23.3)		
Gender										
Male	*	*	*	*	4,352	167,200	16.4	(11.8 - 20.9)		
Female	138	8,900	8.0	(3.1 - 12.9)	7,987	324,000	21.8	(16.6 - 27.0)		
Age (years)										
≥ 65	217	15,800	8.2	(4.3 - 12.2)	12,339	491,200	19.6	(16.0 - 23.3)		
Race/Ethnicity										
White Non-Hispanic	205	13,600	7.8	(3.9 - 11.8)	11,885	322,600	16.8	(14.3 - 19.2)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	6,358	314,100	25.3	(19.9 - 30.7)		
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)		
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)		
Income										
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)		
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)		
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)		
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)		

 $<sup>^{</sup>m 1}$  Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	lassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	217	8.3	(4.3 - 12.3)	12,339	19.4	(15.7 - 23.2)		
Gender								
Male	*	*	*	4,352	16.5	(11.5 - 21.4)		
Female	138	8.0	(3.1 - 12.9)	7,987	22.1	(16.6 - 27.7)		
Race/Ethnicity								
White Non-Hispanic	205	7.9	(3.8 - 12.0)	11,885	16.6	(13.9 - 19.3)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	6,358	24.9	(19.3 - 30.5)		
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)		
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)		
Income								
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)		
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)		
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)		
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)		

 $<sup>^{1}</sup>$  Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 18a. Flu shot within the past 12 months<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	443,200	44.3	(39.1 - 49.5)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	242	198,900	41.8	(33.8 - 49.8)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	404	244,300	46.6	(40.0 - 53.2)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	235	174,400	33.2	(25.0 - 41.5)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	192	116,700	41.5	(32.9 - 50.0)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	219	152,100	78.4	(71.9 - 84.9)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	553	341,800	48.2	(43.1 - 53.4)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	144	84,900	36.2	(26.4 - 46.0)	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	343	223,900	44.9	(38.0 - 51.9)	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	*	*	*	*	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	*	*	*	*	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	276	201,600	45.5	(37.8 - 53.2)	8,548	1,776,100	39.6	(36.3 - 42.8)

<sup>&</sup>lt;sup>1</sup> Defined as having had a flu shot within the past 12 months

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

 <sup>4 95%</sup> confidence interval
 \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 18b. Flu shot within the past 12 months<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	P	Nassau C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	42.6	(37.0 - 48.3)	38,079	41.7	(39.2 - 44.1)
Gender						
Male	242	40.9	(32.5 - 49.4)	14,579	40.0	(36.2 - 43.8)
Female	404	44.2	(36.6 - 51.8)	23,500	43.1	(40.0 - 46.3)
Race/Ethnicity						
White Non-Hispanic	553	46.0	(40.3 - 51.6)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,663	43.7	(39.1 - 48.2)
Some College	*	*	*	9,898	37.8	(33.0 - 42.7)
≥ College Degree	343	43.1	(35.5 - 50.7)	12,436	42.5	(39.1 - 46.0)
Income						
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	*	*	*	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	276	44.4	(36.1 - 52.8)	8,548	39.1	(35.8 - 42.4)

 $<sup>^{\</sup>rm l}$  Defined as having had a flu shot within the past 12 months

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Nassau County and New York State

		Nassa	au County	,		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)		
Gender										
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)		
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)		
Age (years)										
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)		
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)		
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)		
Race/Ethnicity										
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)		
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)		
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)		
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)		
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)		
Income										
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)		
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)		
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)		
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)		

 $<sup>^{\</sup>rm l}$  Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	1	Nassau Co	unty	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	
Total	*	*	*	38,106	1.7	(0.6 - 2.7)	
Gender							
Male	*	*	*	14,586	2.0	(0.1 - 3.9)	
Female	*	*	*	23,520	1.4	(0.5 - 2.3)	
Race/Ethnicity							
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)	
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)	
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)	
Other Non-Hispanic	*	*	*	*	*	*	
Education							
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)	
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)	
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)	
Income							
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)	
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)	
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)	
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)	

 $<sup>^{\</sup>rm I}$  Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	446,900	44.7	(39.5 - 49.9)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	242	202,600	42.6	(34.5 - 50.6)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	404	244,300	46.6	(40.0 - 53.2)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	235	178,100	34.0	(25.6 - 42.3)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	192	116,700	41.5	(32.9 - 50.0)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	219	152,100	78.4	(71.9 - 84.9)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	553	341,800	48.2	(43.1 - 53.4)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	144	88,600	37.8	(27.8 - 47.8)	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	343	223,900	44.9	(38.0 - 51.9)	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	*	*	*	*	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	*	*	*	*	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	276	205,300	46.3	(38.6 - 54.0)	8,548	1,791,200	39.9	(36.6 - 43.2)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest

 <sup>4 95%</sup> confidence interval
 \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	P	Nassau C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	43.0	(37.4 - 48.7)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	242	41.7	(33.2 - 50.2)	14,568	40.4	(36.6 - 44.2)
Female	404	44.2	(36.6 - 51.8)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	553	46.0	(40.3 - 51.6)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,651	44.1	(39.6 - 48.6)
Some College	*	*	*	9,895	38.1	(33.2 - 43.0)
≥ College Degree	343	43.1	(35.5 - 50.7)	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	*	*	*	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	276	45.3	(37.0 - 53.7)	8,548	39.4	(36.1 - 42.7)

 $<sup>^{\</sup>rm I}$  Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 21a. Flu shot among adults aged 65 and older in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	219	152,100	78.4	(71.9 - 84.9)	12,602	1,907,700	75.2	(72.0 - 78.4)		
Gender										
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)		
Female	140	89,400	79.4	(71.5 - 87.2)	8,185	1,134,800	75.6	(71.6 - 79.6)		
Age (years)										
≥ 65	219	152,100	78.4	(71.9 - 84.9)	12,602	1,907,700	75.2	(72.0 - 78.4)		
Race/Ethnicity										
White Non-Hispanic	207	141,400	80.3	(73.9 - 86.6)	12,137	1,487,100	76.0	(73.5 - 78.6)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	6,518	930,800	74.0	(68.9 - 79.2)		
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)		
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)		
Income										
≤ \$24,999	*	*	*	*	4,665	545,300	72.4	(65.6 - 79.3)		
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)		
\$50,000-\$74,999	*	*	*	*	*	*	*	*		
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)		

 $<sup>^{</sup>m 1}$  Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 21b. Flu shot among adults aged 65 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau County			New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	219	78.3	(71.7 - 85.0)	12,602	75.0	(71.5 - 78.5)		
Gender								
Male	*	*	*	4,417	74.4	(68.7 - 80.1)		
Female	140	79.5	(71.7 - 87.2)	8,185	75.5	(71.3 - 79.8)		
Race/Ethnicity								
White Non-Hispanic	207	80.2	(73.5 - 86.8)	12,137	75.9	(73.0 - 78.8)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	6,518	73.3	(67.5 - 79.1)		
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)		
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)		
Income								
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)		
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a flu shot during the past 12 months among adults aged 65 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 22a. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	595	214,600	24.1	(20.3 - 28.0)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	216	94,700	22.1	(16.4 - 27.9)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	379	119,900	26.0	(21.0 - 31.0)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	203	37,900	8.8	(4.6 - 13.1)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	179	44,500	16.5	(10.4 - 22.6)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	213	132,100	70.0	(62.6 - 77.3)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	509	192,400	30.6	(26.2 - 35.0)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	149	77,000	31.7	(22.5 - 40.8)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	131	42,500	21.7	(14.3 - 29.0)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	312	93,100	20.8	(16.1 - 25.6)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	*	*	*	*	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	*	*	*	*	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	*	*	*	*	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	250	64,800	16.8	(12.0 - 21.5)	7,773	648,700	16.3	(14.1 - 18.5)

<sup>&</sup>lt;sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 22b. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	595	22.3	(18.5 - 26.2)	35,303	25.4	(23.2 - 27.6)		
Gender								
Male	216	22.3	(16.2 - 28.3)	13,052	25.8	(22.3 - 29.4)		
Female	379	22.3	(17.5 - 27.2)	22,251	25.1	(22.4 - 27.8)		
Race/Ethnicity								
White Non-Hispanic	509	28.7	(24.1 - 33.3)	33,225	27.7	(25.9 - 29.5)		
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)		
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)		
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)		
Education								
≤ High School	149	29.7	(20.1 - 39.2)	14,700	29.5	(25.6 - 33.5)		
Some College	131	20.7	(13.3 - 28.2)	9,151	26.2	(21.0 - 31.3)		
≥ College Degree	312	18.9	(14.2 - 23.6)	11,375	21.0	(18.3 - 23.8)		
Income								
≤ \$24,999	*	*	*	9,082	32.3	(26.9 - 37.8)		
\$25,000-\$49,999	*	*	*	9,033	29.0	(23.0 - 35.0)		
\$50,000-\$74,999	*	*	*	5,320	21.2	(16.8 - 25.7)		
≥ \$75,000	250	15.6	(10.8 - 20.3)	7,773	16.2	(13.9 - 18.5)		

 $<sup>^{\</sup>rm 1}$  Defined as having ever had a pneumonia shot or pneumococcal vaccine

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{\</sup>rm 3}$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	213	132,100	70.0	(62.6 - 77.3)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	140	77,300	68.6	(59.2 - 78.0)	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	213	132,100	70.0	(62.6 - 77.3)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	201	123,300	72.2	(64.9 - 79.5)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	*	*	*	*	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

 $<sup>^{</sup>m 1}$  Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	Nassau C	ounty	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	213	70.3	(63.0 - 77.7)	12,181	64.7	(60.4 - 69.1)		
Gender								
Male	*	*	*	4,180	61.5	(54.3 - 68.6)		
Female	140	68.8	(59.5 - 78.1)	8,001	67.6	(62.4 - 72.9)		
Race/Ethnicity								
White Non-Hispanic	201	72.3	(64.9 - 79.7)	11,735	67.1	(63.2 - 70.9)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	6,345	64.2	(57.4 - 71.0)		
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)		
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)		
Income								
≤ \$24,999	*	*	*	4,533	66.0	(57.0 - 74.9)		
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	*	*	*		

<sup>&</sup>lt;sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 24a. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	y		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	333	352,100	97.8	(96.3 - 99.4)	19,491	4,328,800	91.8	(89.7 - 94.0)		
Gender										
Male	-	-	-	-	-	-	-	-		
Female	333	352,100	97.8	(96.3 - 99.4)	19,491	4,328,800	91.8	(89.7 - 94.0)		
Age (years)										
40-49	84	111,400	98.9	(96.7 - 100)	3,851	1,287,500	85.0	(79.7 - 90.3)		
50-64	109	131,300	97.5	(94.3 - 100)	7,452	1,641,700	96.0	(94.1 - 97.9)		
≥ 65	140	109,400	97.1	(94.8 - 99.5)	8,188	1,399,600	94.1	(90.5 - 97.6)		
Race/Ethnicity										
White Non-Hispanic	295	284,100	97.6	(95.8 - 99.5)	18,548	2,976,100	93.9	(92.7 - 95.1)		
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)		
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	84	93,500	98.1	(95.9 - 100)	8,266	1,698,700	91.0	(87.4 - 94.5)		
Some College	83	83,600	98.1	(95.1 - 100)	5,322	1,055,500	93.6	(90.2 - 97.0)		
≥ College Degree	165	174,000	97.5	(95.0 - 100)	5,872	1,560,700	91.6	(87.5 - 95.6)		
Income										
≤ \$24,999	*	*	*	*	5,691	1,023,400	93.5	(89.7 - 97.4)		
\$25,000-\$49,999	55	57,400	94.6	(87.6 - 100)	4,857	860,300	89.5	(83.6 - 95.5)		
\$50,000-\$74,999	56	61,900	99.2	(97.5 - 100)	2,648	547,100	94.6	(92.2 - 96.9)		
≥ \$75,000	129	148,500	98.3	(96.2 - 100)	3,582	1,185,100	92.2	(88.2 - 96.1)		

 $<sup>^{1}\,\</sup>mathrm{Defined}$  as having ever had a mammogram among women aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 24b. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	333	98.0	(96.6 - 99.4)	19,491	91.1	(88.7 - 93.6)		
Gender								
Male	-	-	-	-	-	-		
Female	333	98.0	(96.6 - 99.4)	19,491	91.1	(88.7 - 93.6)		
Race/Ethnicity								
White Non-Hispanic	295	97.8	(96.2 - 99.5)	18,548	93.3	(92.0 - 94.7)		
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)		
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	84	98.2	(96.0 - 100)	8,266	90.2	(86.1 - 94.2)		
Some College	83	98.1	(95.2 - 100)	5,322	92.9	(88.9 - 96.9)		
≥ College Degree	165	97.9	(95.7 - 100)	5,872	90.9	(86.6 - 95.3)		
Income								
≤ \$24,999	*	*	*	5,691	93.0	(88.5 - 97.4)		
\$25,000-\$49,999	55	95.1	(88.8 - 100)	4,857	88.8	(82.5 - 95.1)		
\$50,000-\$74,999	56	99.2	(97.7 - 100)	2,648	94.3	(91.7 - 96.8)		
≥ \$75,000	129	98.5	(96.6 - 100)	3,582	91.2	(86.8 - 95.6)		

 $<sup>^{\</sup>rm 1}$  Defined as having ever had a mammogram among women aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	331	304,300	85.2	(80.6 - 89.8)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	_	-	-	-
Female	331	304,300	85.2	(80.6 - 89.8)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	109	122,400	90.9	(85.5 - 96.4)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	139	94,600	84.4	(77.2 - 91.6)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	294	252,400	86.9	(82.3 - 91.4)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	82	78,800	85.2	(77.5 - 92.9)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	83	73,200	85.9	(76.2 - 95.6)	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	165	151,200	84.8	(77.8 - 91.8)	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	*	*	*	*	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	*	*	*	*	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	129	130,600	86.4	(79.1 - 93.7)	3,572	1,025,900	80.0	(75.2 - 84.8)

 $<sup>^{\</sup>rm 1}$  Defined as having had a mammogram within the past 2 years among women aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 25b. Had mammogram within the past 2 years among women aged 40 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	ounty	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	331	85.3	(80.7 - 90.0)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	331	85.3	(80.7 - 90.0)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	294	86.9	(82.3 - 91.5)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	82	84.7	(76.7 - 92.7)	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	165	85.7	(79.1 - 92.3)	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	*	*	*	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	129	86.6	(79.3 - 93.9)	3,572	79.0	(73.8 - 84.2)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a mammogram within the past 2 years among women aged 40 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	248	217,000	88.0	(83.5 - 92.5)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	248	217,000	88.0	(83.5 - 92.5)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	109	122,400	90.9	(85.5 - 96.4)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	139	94,600	84.4	(77.2 - 91.6)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	225	186,100	89.3	(85.1 - 93.5)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	69	58,900	85.2	(77.0 - 93.4)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	58	50,900	96.6	(92.4 - 100)	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	120	106,100	85.7	(78.3 - 93.2)	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	*	*	*	*	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	*	*	*	*	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	83	83,400	92.9	(87.5 - 98.3)	2,303	601,300	84.6	(79.3 - 89.9)

<sup>&</sup>lt;sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 26b. Had mammogram within the past 2 years among women aged 50 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	ľ	Nassau C	County	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	248	88.2	(83.8 - 92.6)	15,534	83.1	(80.3 - 85.8)
Gender						
Male	-	-	-	-	-	-
Female	248	88.2	(83.8 - 92.6)	15,534	83.1	(80.3 - 85.8)
Race/Ethnicity						
White Non-Hispanic	225	89.4	(85.2 - 93.6)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	69	85.3	(77.2 - 93.4)	7,026	83.6	(80.1 - 87.0)
Some College	58	97.0	(93.4 - 100)	4,124	87.3	(84.1 - 90.5)
≥ College Degree	120	86.0	(78.8 - 93.3)	4,356	79.9	(73.9 - 86.0)
Income						
≤ \$24,999	*	*	*	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	*	*	*	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	83	92.9	(87.5 - 98.4)	2,303	83.9	(77.9 - 89.9)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a mammogram within the past 2 years among women aged 50 and older

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 27a. Ever had a Pap test among women<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	401	476,700	91.5	(85.0 - 97.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	401	476,700	91.5	(85.0 - 97.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	79	92,000	100	(100 - 100)	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	69	111,500	100	(100 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	82	87,600	100	(100 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	139	110,300	98.4	(96.5 - 100)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	345	364,200	95.2	(90.5 - 100)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	*	*	*	*	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	206	250,800	98.2	(95.0 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	*	*	*	*	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	63	77,700	100	(100 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	65	85,300	100	(100 - 100)	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	163	202,500	94.7	(87.3 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

 $<sup>^{\</sup>rm I}$  Defined as having ever had a Pap test among women

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 27b. Ever had a Pap test among women<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	Nassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	401	88.7	(80.5 - 97.0)	23,381	91.8	(89.6 - 94.0)		
Gender								
Male	-	-	-	-	-	-		
Female	401	88.7	(80.5 - 97.0)	23,381	91.8	(89.6 - 94.0)		
Race/Ethnicity								
White Non-Hispanic	345	93.3	(86.5 - 100)	21,976	94.0	(91.8 - 96.2)		
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)		
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	9,526	87.8	(83.2 - 92.4)		
Some College	*	*	*	6,462	93.1	(89.7 - 96.4)		
≥ College Degree	206	97.5	(93.0 - 100)	7,345	94.9	(92.0 - 97.7)		
Income								
≤ \$24,999	*	*	*	6,526	93.3	(90.1 - 96.5)		
\$25,000-\$49,999	63	100	(100 - 100)	5,787	91.1	(85.7 - 96.5)		
\$50,000-\$74,999	65	100	(100 - 100)	3,310	96.0	(91.9 - 100)		
≥ \$75,000	*	*	*	4,570	95.0	(91.9 - 98.1)		

<sup>&</sup>lt;sup>1</sup> Defined as having ever had a Pap test among women

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 28a. Had a Pap test within the past 3 years among women<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	е
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	396	412,800	79.7	(72.9 - 86.5)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	396	412,800	79.7	(72.9 - 86.5)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	79	85,000	92.4	(82.8 - 100)	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	69	102,100	91.7	(83.7 - 99.6)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	*	*	*	*	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	135	83,300	76.1	(68.6 - 83.7)	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	340	319,900	84.3	(78.9 - 89.8)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	*	*	*	*	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	203	224,100	88.3	(82.8 - 93.7)	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	*	*	*	*	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	65	79,500	93.2	(88.1 - 98.2)	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	163	190,900	89.2	(81.1 - 97.3)	4,555	1,810,200	89.0	(85.7 - 92.2)

 $<sup>^{\</sup>rm I}$  Defined as having had a Pap test within the past 3 years among women

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 28b. Had a Pap test within the past 3 years among women<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	1	lassau C	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	396	79.1	(71.0 - 87.2)	23,109	82.7	(80.2 - 85.1)
Gender						
Male	-	-	-	-	-	-
Female	396	79.1	(71.0 - 87.2)	23,109	82.7	(80.2 - 85.1)
Race/Ethnicity						
White Non-Hispanic	340	84.2	(77.3 - 91.2)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	9,384	76.4	(71.3 - 81.5)
Some College	*	*	*	6,401	84.2	(80.5 - 88.0)
≥ College Degree	203	89.7	(84.0 - 95.4)	7,281	87.5	(84.3 - 90.6)
Income						
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	*	*	*	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	65	94.2	(89.9 - 98.6)	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	*	*	*	4,555	88.9	(85.5 - 92.3)

 $<sup>^{\</sup>rm 1}$  Defined as having had a Pap test within the past 3 years among women

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 29a. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	195	271,400	85.9	(79.8 - 92.1)	12,033	3,176,500	73.8	(69.7 - 77.8)		
Gender										
Male	195	271,400	85.9	(79.8 - 92.1)	12,033	3,176,500	73.8	(69.7 - 77.8)		
Female	-	-	-	-	-	-	-	-		
Age (years)										
40-44	*	*	*	*	*	*	*	*		
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)		
55-64	53	73,700	92.4	(83.7 - 100)	3,504	865,600	87.1	(82.5 - 91.7)		
≥ 65	79	77,700	95.4	(91.7 - 99.1)	4,348	953,900	92.9	(91.0 - 94.9)		
Race/Ethnicity										
White Non-Hispanic	174	210,800	87.3	(81.2 - 93.3)	11,411	2,452,100	80.3	(77.7 - 82.9)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)		
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)		
≥ College Degree	111	142,600	86.7	(78.4 - 94.9)	4,302	1,568,000	80.5	(74.0 - 87.0)		
Income										
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)		
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)		
\$50,000-\$74,999	*	*	*	*	*	*	*	*		
≥ \$75,000	94	151,900	88.4	(80.3 - 96.5)	3,309	1,351,100	78.0	(72.6 - 83.5)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a digital rectal exam among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 29b. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	lassau C	ounty	N	New York State			
,	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total 19	95	85.5	(79.1 - 92.0)	12,033	73.7	(69.6 - 77.8)		
Gender								
Male 19	95	85.5	(79.1 - 92.0)	12,033	73.7	(69.6 - 77.8)		
Female	-	-	-	-	-	-		
Race/Ethnicity								
White Non-Hispanic 1	74	86.3	(79.5 - 93.1)	11,411	79.9	(77.2 - 82.5)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)		
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)		
≥ College Degree 1	11	85.3	(75.8 - 94.8)	4,302	80.3	(73.8 - 86.8)		
Income								
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)		
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	94	87.0	(77.7 - 96.4)	3,309	77.5	(72.0 - 83.0)		

 $<sup>^{\</sup>rm I}$  Defined as having ever had a digital rectal exam among men aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	194	210,000	67.1	(57.6 - 76.5)	11,923	2,380,300	55.8	(51.8 - 59.8)		
Gender										
Male	194	210,000	67.1	(57.6 - 76.5)	11,923	2,380,300	55.8	(51.8 - 59.8)		
Female	-	-	-	-	-	-	-	-		
Age (years)										
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)		
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)		
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)		
≥ 65	79	66,100	81.1	(71.2 - 91.0)	4,304	809,900	79.7	(75.0 - 84.3)		
Race/Ethnicity										
White Non-Hispanic	173	169,700	71.0	(63.0 - 79.0)	11,305	1,875,600	62.2	(59.2 - 65.2)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)		
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)		
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)		
Income										
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)		
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)		
\$50,000-\$74,999	*	*	*	*	*	*	*	*		
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Nassau County and New York State - Age-Adjusted 2

	1	Nassau C	ounty	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	*	*	*	11,923	55.9	(51.7 - 60.0)		
Gender								
Male	*	*	*	11,923	55.9	(51.7 - 60.0)		
Female	-	-	-	-	-	-		
Race/Ethnicity								
White Non-Hispanic	173	70.1	(61.7 - 78.6)	11,305	62.0	(58.9 - 65.1)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)		
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)		
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)		
Income								
≤ \$24,999	*	*	*	*	*	*		
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	188	224,900	73.5	(65.4 - 81.6)	11,557	2,822,600	67.6	(63.9 - 71.2)
Gender								
Male	188	224,900	73.5	(65.4 - 81.6)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	74	72,500	94.8	(88.8 - 100)	4,215	905,400	90.4	(87.1 - 93.6)
Race/Ethnicity								
White Non-Hispanic	167	168,500	72.1	(63.5 - 80.7)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a prostate specific antigen test among men aged 40 and older

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	P	Nassau C	ounty	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	188	71.9	(63.3 - 80.6)	11,557	67.4	(63.7 - 71.1)
Gender						
Male	188	71.9	(63.3 - 80.6)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	167	69.6	(60.4 - 78.9)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

<sup>&</sup>lt;sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	187	208,200	68.2	(59.8 - 76.7)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	187	208,200	68.2	(59.8 - 76.7)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	74	68,800	89.9	(82.3 - 97.5)	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	166	151,700	65.2	(56.4 - 74.1)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75 <b>,</b> 000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

 $<sup>^{</sup>m 1}$  Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	187	66.7	(57.8 - 75.7)	11,437	59.8	(55.8 - 63.9)		
Gender								
Male	187	66.7	(57.8 - 75.7)	11,437	59.8	(55.8 - 63.9)		
Female	-	-	-	-	-	-		
Race/Ethnicity								
White Non-Hispanic	166	62.9	(53.6 - 72.2)	10,836	60.1	(56.9 - 63.4)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)		
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)		
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)		
Income								
≤ \$24,999	*	*	*	*	*	*		
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 33a. Prostate cancer among men aged 40 and older in Nassau County and New York State

		Nassa	au County	′		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	197	21,600	6.8	(3.3 - 10.2)	12,133	178,200	4.1	(3.0 - 5.2)		
Gender										
Male	197	21,600	6.8	(3.3 - 10.2)	12,133	178,200	4.1	(3.0 - 5.2)		
Female	-	-	-	-	-	-	-	-		
Age (years)										
40-44	*	*	*	*	*	*	*	*		
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)		
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)		
≥ 65	79	15,200	18.6	(9.4 - 27.9)	4,388	140,400	13.6	(9.7 - 17.4)		
Race/Ethnicity										
White Non-Hispanic	175	20,000	8.3	(4.0 - 12.6)	11,502	143,700	4.7	(3.5 - 5.8)		
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)		
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)		
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)		
≥ College Degree	111	13,400	8.2	(2.7 - 13.6)	4,336	92,500	4.7	(3.0 - 6.5)		
Income										
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)		
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)		
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)		
≥ \$75,000	94	13,200	7.7	(2.3 - 13.1)	3,322	56,500	3.2	(1.7 - 4.8)		

<sup>&</sup>lt;sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 33b. Prostate cancer among men aged 40 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	197	6.9	(3.5 - 10.3)	12,133	4.5	(3.2 - 5.7)		
Gender								
Male	197	6.9	(3.5 - 10.3)	12,133	4.5	(3.2 - 5.7)		
Female	-	-	-	-	-	-		
Race/Ethnicity								
White Non-Hispanic	175	8.3	(4.2 - 12.5)	11,502	5.1	(3.8 - 6.4)		
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)		
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)		
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)		
≥ College Degree	111	8.0	(2.8 - 13.2)	4,336	5.2	(3.1 - 7.2)		
Income								
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)		
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)		
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)		
≥ \$75,000	94	7.4	(2.3 - 12.5)	3,322	3.5	(1.6 - 5.4)		

<sup>&</sup>lt;sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 34a. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	408	187,300	39.5	(33.6 - 45.4)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	157	85,800	38.3	(29.0 - 47.5)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	251	101,500	40.7	(33.3 - 48.1)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	190	89,700	31.9	(23.7 - 40.2)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	218	97,700	50.6	(43.0 - 58.1)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	370	156,500	40.1	(34.1 - 46.0)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	209	89,200	39.2	(31.4 - 46.9)	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	*	*	*	*	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	*	*	*	*	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	156	70,200	35.3	(26.3 - 44.4)	4,647	657,800	37.3	(33.0 - 41.6)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever used a home blood stool test among adults aged 50 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 34b. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	408	40.0	(34.1 - 45.8)	25,322	38.1	(35.5 - 40.8)		
Gender								
Male	157	39.2	(30.2 - 48.3)	9,571	39.4	(35.3 - 43.5)		
Female	251	40.7	(33.2 - 48.1)	15,751	36.9	(33.5 - 40.3)		
Race/Ethnicity								
White Non-Hispanic	370	40.4	(34.4 - 46.4)	24,213	38.9	(36.7 - 41.1)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	11,146	35.1	(30.3 - 39.9)		
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)		
≥ College Degree	209	40.0	(32.1 - 47.8)	7,851	41.9	(37.9 - 46.0)		
Income								
≤ \$24,999	*	*	*	7,222	35.9	(29.5 - 42.4)		
\$25,000-\$49,999	*	*	*	6,609	40.2	(34.8 - 45.6)		
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)		
≥ \$75,000	156	35.5	(26.4 - 44.7)	4,647	37.5	(32.9 - 42.2)		

 $<sup>^{\</sup>rm I}$  Defined as having ever used a home blood stool test among adults aged 50 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	405	71,700	15.2	(11.0 - 19.4)	25,074	863,300	14.4	(12.6 - 16.2)		
Gender										
Male	156	39,000	17.5	(10.3 - 24.7)	9,508	444,400	16.0	(13.2 - 18.9)		
Female	249	32,700	13.2	(8.4 - 17.9)	15,566	418,900	13.1	(10.8 - 15.4)		
Age (years)										
50-64	189	27,700	9.9	(4.7 - 15.1)	12,759	454,400	13.0	(10.6 - 15.5)		
≥ 65	216	44,000	23.0	(16.2 - 29.7)	12,299	406,400	16.4	(13.7 - 19.0)		
Race/Ethnicity										
White Non-Hispanic	367	60,200	15.5	(11.3 - 19.7)	23,971	634,300	14.3	(12.8 - 15.7)		
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)		
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	109	15,600	12.3	(6.1 - 18.6)	11,026	288,600	12.3	(9.4 - 15.3)		
Some College	*	*	*	*	6,208	212,500	16.4	(12.5 - 20.4)		
≥ College Degree	207	35,600	15.8	(10.0 - 21.5)	7,778	352,400	15.2	(12.5 - 17.9)		
Income										
≤ \$24,999	*	*	*	*	7,157	196,300	13.9	(10.1 - 17.6)		
\$25,000-\$49,999	*	*	*	*	6,562	221,500	19.2	(14.1 - 24.3)		
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)		
≥ \$75,000	155	20,300	10.3	(5.8 - 14.8)	4,623	205,100	11.7	(8.8 - 14.6)		

 $<sup>^{</sup>m 1}$  Defined as having used a home blood stool test within the past year among adults aged 50 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

4 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a

numerator less than 10)

Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Nassau County and New York State - Age-Adjusted 2

	r	Nassau C	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	405	15.6	(11.4 - 19.8)	25,074	14.6	(12.6 - 16.7)
Gender						
Male	156	18.3	(11.3 - 25.2)	9,508	16.4	(13.2 - 19.6)
Female	249	13.0	(8.3 - 17.7)	15,566	12.9	(10.4 - 15.4)
Race/Ethnicity						
White Non-Hispanic	367	16.1	(11.7 - 20.5)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	109	13.1	(6.5 - 19.6)	11,026	12.7	(9.2 - 16.1)
Some College	86	16.5	(6.7 - 26.3)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	207	16.3	(10.3 - 22.3)	7,778	15.3	(12.4 - 18.3)
Income						
≤ \$24,999	*	*	*	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	*	*	*	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	155	10.5	(5.8 - 15.1)	4,623	11.8	(8.6 - 15.0)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having used a home blood stool test within the past year among adults aged 50 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	405	98,800	21.0	(16.1 - 25.8)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	156	47,100	21.1	(13.5 - 28.7)	9,508	593,100	21.4	(18.3 - 24.5)
Female	249	51,800	20.9	(14.8 - 27.0)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	189	42,500	15.2	(9.0 - 21.4)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	216	56,300	29.4	(22.1 - 36.7)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	367	81,400	21.0	(16.1 - 25.8)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	109	26,200	20.6	(12.2 - 29.1)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	*	*	*	*	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	207	48,900	21.7	(14.9 - 28.5)	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	*	*	*	*	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	*	*	*	*	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	155	34,500	17.5	(10.9 - 24.0)	4,623	294,900	16.8	(13.7 - 20.0)

 $<sup>^{</sup>m 1}$  Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

 $<sup>^{2}\,\</sup>mathrm{Sample}$  size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	assau C	ounty	N	New York State			
n	1 <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total 40	)5	21.2	(16.4 - 26.0)	25,074	20.5	(18.2 - 22.8)		
Gender								
Male 15	6	21.8	(14.5 - 29.2)	9,508	21.7	(18.3 - 25.1)		
Female 24	19	20.6	(14.5 - 26.8)	15,566	19.4	(16.3 - 22.5)		
Race/Ethnicity								
White Non-Hispanic 36	57	21.3	(16.4 - 26.3)	23,971	19.7	(18.0 - 21.3)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School 10	)9	21.1	(12.6 - 29.6)	11,026	19.5	(15.4 - 23.7)		
Some College	*	*	*	6,208	20.4	(16.1 - 24.7)		
≥ College Degree 20	)7	22.2	(15.2 - 29.2)	7,778	21.4	(18.1 - 24.7)		
Income								
≤ \$24,999	*	*	*	7,157	18.9	(14.2 - 23.6)		
\$25,000-\$49,999	*	*	*	6,562	24.8	(19.3 - 30.3)		
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)		
≥ \$75,000 15	55	17.8	(11.1 - 24.4)	4,623	16.9	(13.4 - 20.4)		

 $<sup>^{</sup>m 1}$  Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	409	335,300	70.7	(64.9 - 76.6)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	158	149,200	66.3	(56.5 - 76.1)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	251	186,000	74.7	(68.2 - 81.3)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	136	122,000	72.7	(63.3 - 82.0)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	218	154,800	80.2	(74.1 - 86.3)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	371	282,600	72.3	(66.5 - 78.0)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	210	174,200	76.2	(69.2 - 83.2)	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	*	*	*	*	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	*	*	*	*	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	72	78,400	86.7	(77.9 - 95.5)	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Nassau County and New York State - Age-Adjusted 2

	r	lassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	409	71.0	(65.2 - 76.7)	25,357	66.6	(63.9 - 69.3)		
Gender								
Male	158	67.7	(58.4 - 76.9)	9,582	68.5	(64.5 - 72.5)		
Female	251	74.2	(67.5 - 80.9)	15,775	64.8	(61.2 - 68.4)		
Race/Ethnicity								
White Non-Hispanic	371	72.6	(66.9 - 78.3)	24,248	68.7	(66.5 - 70.8)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	11,121	61.3	(56.4 - 66.2)		
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)		
≥ College Degree	210	76.9	(70.0 - 83.8)	7,884	74.0	(70.2 - 77.9)		
Income								
≤ \$24,999	*	*	*	7,220	63.0	(56.7 - 69.3)		
\$25,000-\$49,999	*	*	*	6,619	59.7	(53.8 - 65.6)		
\$50,000-\$74,999	72	86.7	(78.0 - 95.3)	3,514	74.4	(68.3 - 80.5)		
≥ \$75,000	156	67.3	(57.5 - 77.1)	4,665	73.6	(69.3 - 78.0)		

 $<sup>^{\</sup>rm 1}$  Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	407	329,500	69.7	(63.8 - 75.6)	25,169	3,855,800	64.6	(62.1 - 67.1)		
Gender										
Male	157	148,000	65.9	(56.1 - 75.7)	9,521	1,835,400	66.4	(62.6 - 70.1)		
Female	250	181,500	73.2	(66.5 - 79.8)	15,648	2,020,400	63.1	(59.8 - 66.4)		
Age (years)										
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)		
55-64	135	120,500	72.0	(62.6 - 81.4)	8,652	1,455,100	69.4	(65.7 - 73.2)		
≥ 65	217	152,800	79.5	(73.4 - 85.7)	12,331	1,695,700	69.5	(65.9 - 73.1)		
Race/Ethnicity										
White Non-Hispanic	370	277,800	71.1	(65.3 - 76.9)	24,069	2,981,500	66.6	(64.6 - 68.6)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	11,017	1,360,400	59.0	(54.5 - 63.5)		
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)		
≥ College Degree	210	171,400	75.0	(67.9 - 82.1)	7,841	1,682,500	71.7	(68.0 - 75.4)		
Income										
≤ \$24,999	*	*	*	*	7,163	858,900	61.0	(55.1 - 67.0)		
\$25,000-\$49,999	*	*	*	*	6,584	664,900	57.7	(52.2 - 63.2)		
\$50,000-\$74,999	72	77,700	86.0	(77.2 - 94.9)	3,501	602,600	73.1	(67.7 - 78.5)		
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)		

 $<sup>^{</sup>m 1}$  Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	P	Nassau C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	407	69.9	(64.2 - 75.7)	25,169	64.6	(61.9 - 67.3)
Gender						
Male	157	67.3	(58.0 - 76.5)	9,521	66.9	(62.9 - 70.9)
Female	250	72.6	(65.8 - 79.4)	15,648	62.4	(58.9 - 66.0)
Race/Ethnicity						
White Non-Hispanic	370	71.5	(65.7 - 77.2)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	210	75.5	(68.6 - 82.5)	7,841	72.1	(68.2 - 75.9)
Income						
≤ \$24,999	*	*	*	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	*	*	*	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	72	85.8	(77.0 - 94.6)	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	156	66.1	(56.4 - 75.9)	4,655	71.8	(67.4 - 76.2)

 $<sup>^{</sup>m I}$  Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	403	339,600	72.4	(66.5 - 78.2)	24,792	4,049,200	69.1	(66.7 - 71.6)		
Gender										
Male	156	153,600	68.6	(58.7 - 78.4)	9,418	1,945,200	70.7	(67.0 - 74.3)		
Female	247	186,000	75.9	(69.4 - 82.3)	15,374	2,104,100	67.8	(64.6 - 71.1)		
Age (years)										
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)		
55-64	131	120,700	73.6	(64.2 - 83.0)	8,366	1,487,600	73.7	(70.0 - 77.4)		
≥ 65	217	160,900	83.8	(78.3 - 89.3)	12,294	1,810,500	74.0	(70.6 - 77.4)		
Race/Ethnicity										
White Non-Hispanic	366	287,900	74.4	(68.7 - 80.0)	23,726	3,091,400	70.4	(68.4 - 72.4)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	10,872	1,461,600	64.2	(59.8 - 68.7)		
Some College	*	*	*	*	6,170	873,300	68.4	(63.9 - 72.8)		
≥ College Degree	208	174,700	77.3	(70.3 - 84.2)	7,708	1,709,000	74.6	(70.9 - 78.3)		
Income										
≤ \$24,999	*	*	*	*	7,092	923,300	66.0	(60.1 - 71.9)		
\$25,000-\$49,999	*	*	*	*	6,526	730,000	64.2	(58.9 - 69.6)		
\$50,000-\$74,999	72	79,000	87.4	(78.9 - 95.9)	3,467	621,500	76.6	(71.4 - 81.9)		
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)		

<sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older <sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	lassau C	ounty	N	New York State		
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	
Total	403	72.7	(67.0 - 78.4)	24,792	69.3	(66.7 - 71.9)	
Gender							
Male	156	70.1	(60.8 - 79.3)	9,418	71.5	(67.6 - 75.4)	
Female	247	75.3	(68.7 - 81.9)	15,374	67.2	(63.7 - 70.7)	
Race/Ethnicity							
White Non-Hispanic	366	74.8	(69.1 - 80.4)	23,726	70.3	(68.1 - 72.4)	
Black Non-Hispanic	*	*	*	*	*	*	
Hispanic	*	*	*	*	*	*	
Other Non-Hispanic	*	*	*	*	*	*	
Education							
≤ High School	*	*	*	10,872	64.5	(59.8 - 69.3)	
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)	
≥ College Degree	208	77.7	(70.9 - 84.6)	7,708	75.0	(71.2 - 78.8)	
Income							
≤ \$24,999	*	*	*	7,092	66.4	(60.2 - 72.6)	
\$25,000-\$49,999	*	*	*	6,526	63.8	(58.2 - 69.5)	
\$50,000-\$74,999	72	87.5	(79.3 - 95.8)	3,467	76.7	(70.6 - 82.7)	
≥ \$75,000	156	68.5	(58.7 - 78.3)	4,615	74.4	(70.2 - 78.6)	

 $<sup>^{1}</sup>$  Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 40a. High blood pressure among adults in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	647	278,200	27.8	(23.3 - 32.3)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	243	147,300	30.8	(23.3 - 38.3)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	404	131,000	25.0	(19.9 - 30.1)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	*	*	*	*	4,984	426,200	14.8	(11.6 - 18.1)
45-54	115	53,200	24.5	(15.3 - 33.7)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	*	*	*	*	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	219	104,200	53.8	(46.3 - 61.3)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	554	184,000	25.9	(21.9 - 30.0)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	155	86,000	32.6	(23.2 - 42.0)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	146	65,400	27.5	(18.6 - 36.5)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	343	125,200	25.2	(18.9 - 31.5)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	*	*	*	*	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	*	*	*	*	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	275	97,500	22.1	(15.3 - 29.0)	8,560	942,500	21.0	(18.4 - 23.6)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent has high blood pressure

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 40b. High blood pressure among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	647	24.5	(20.0 - 29.0)	38,125	25.7	(23.8 - 27.5)		
Gender								
Male	243	28.7	(21.0 - 36.4)	14,611	27.7	(24.7 - 30.7)		
Female	404	20.6	(15.9 - 25.3)	23,514	23.8	(21.5 - 26.0)		
Race/Ethnicity								
White Non-Hispanic	554	23.1	(19.2 - 27.1)	35,814	27.2	(25.6 - 28.7)		
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)		
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)		
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)		
Education								
≤ High School	155	28.9	(19.5 - 38.2)	15,688	32.1	(28.4 - 35.8)		
Some College	146	23.7	(15.4 - 32.0)	9,906	22.8	(19.7 - 25.9)		
≥ College Degree	343	22.4	(15.8 - 29.0)	12,452	21.2	(18.6 - 23.8)		
Income								
≤ \$24,999	*	*	*	9,640	31.8	(26.7 - 37.0)		
\$25,000-\$49,999	*	*	*	9,657	26.9	(22.8 - 30.9)		
\$50,000-\$74,999	109	26.8	(17.5 - 36.1)	5,773	24.8	(20.4 - 29.2)		
≥ \$75,000	275	20.0	(12.7 - 27.2)	8,560	19.9	(17.3 - 22.5)		

 $<sup>^{\</sup>rm 1}$  Defined as having ever been told by a health professional that respondent has high blood pressure

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 41a. Blood pressure medication among adults with high blood pressure in Nassau County and New York State

		Nassa	au Count	y		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	219	223,200	80.2	(73.1 - 87.3)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	128	109,300	83.5	(75.4 - 91.5)	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	55	60,700	88.5	(80.0 - 97.1)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	117	96,800	92.9	(87.5 - 98.3)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	183	148,300	80.6	(73.6 - 87.6)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

 $<sup>^{\</sup>mathrm{1}}$  Defined as currently taking medicine for high blood pressure among adults with high blood pressure

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)
<sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 41b. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	219	78.7	(70.8 - 86.6)	15,288	78.5	(75.2 - 81.7)		
Gender								
Male	*	*	*	6,080	73.3	(67.9 - 78.7)		
Female	128	81.3	(72.4 - 90.3)	9,208	84.0	(80.7 - 87.2)		
Race/Ethnicity								
White Non-Hispanic	183	78.6	(70.7 - 86.5)	14,442	79.0	(76.5 - 81.6)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)		
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)		
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)		
Income								
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)		
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)		
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)		
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)		

 $<sup>^{</sup>m 1}$  Defined as currently taking medicine for high blood pressure among adults with high blood pressure

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 42a. Ever had blood cholesterol checked among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	638	852,700	87.7	(83.7 - 91.8)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	240	407,000	85.9	(79.7 - 92.1)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	398	445,700	89.5	(84.2 - 94.8)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	106	158,400	89.2	(81.3 - 97.0)	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	114	206,000	95.7	(91.2 - 100)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	136	161,900	96.8	(93.5 - 100)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	218	189,800	98.4	(96.8 - 99.9)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	546	635,100	91.7	(88.2 - 95.3)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	145	204,900	86.6	(77.9 - 95.4)	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	339	448,500	92.0	(87.6 - 96.3)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	88	114,100	88.9	(80.5 - 97.3)	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	108	145,400	91.7	(84.3 - 99.2)	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	274	392,800	90.2	(84.7 - 95.7)	8,460	4,019,300	92.2	(90.8 - 93.6)

<sup>&</sup>lt;sup>1</sup> Defined as ever having blood cholesterol checked

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 42b. Ever had blood cholesterol checked among adults<sup>1</sup> in Nassau County and New York
State - Age-Adjusted<sup>2</sup>

	r	lassau C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	638	85.7	(80.9 - 90.5)	37,450	79.9	(77.3 - 82.5)
Gender						
Male	240	84.1	(77.2 - 91.0)	14,389	78.0	(73.9 - 82.1)
Female	398	87.3	(80.6 - 94.0)	23,061	81.6	(78.4 - 84.8)
Race/Ethnicity						
White Non-Hispanic	546	89.9	(85.3 - 94.5)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,342	70.2	(65.2 - 75.1)
Some College	*	*	*	9,732	76.7	(71.1 - 82.3)
≥ College Degree	339	90.4	(85.2 - 95.6)	12,296	90.9	(88.6 - 93.2)
Income						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	*	*	*	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	108	89.6	(79.8 - 99.3)	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	274	88.4	(81.7 - 95.1)	8,460	91.7	(90.1 - 93.2)

<sup>&</sup>lt;sup>1</sup> Defined as ever having blood cholesterol checked

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 43a. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	y		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	633	823,100	85.4	(81.1 - 89.6)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	239	391,500	82.7	(76.2 - 89.3)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	394	431,600	87.9	(82.4 - 93.4)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	105	147,300	83.4	(73.9 - 92.8)	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	112	195,200	93.1	(87.5 - 98.6)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	136	158,700	94.9	(91.0 - 98.9)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	216	188,100	98.0	(96.4 - 99.7)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	541	605,500	88.4	(84.4 - 92.4)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	144	199,500	85.0	(76.0 - 94.0)	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	337	429,100	88.8	(84.0 - 93.7)	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	88	113,600	88.6	(80.1 - 97.0)	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	108	138,800	87.6	(78.8 - 96.3)	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	271	372,000	86.7	(80.7 - 92.7)	8,435	3,872,400	89.2	(87.5 - 90.9)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had blood cholesterol checked within the past 5 years

 $<sup>^{2}\ \</sup>mathrm{Sample}\ \mathrm{size}\ \mathrm{from}\ \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 43b. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	633	83.2	(78.2 - 88.1)	37,201	77.3	(74.7 - 79.9)		
Gender								
Male	239	80.5	(73.2 - 87.8)	14,307	74.8	(70.7 - 79.0)		
Female	394	85.8	(78.9 - 92.6)	22,894	79.5	(76.3 - 82.7)		
Race/Ethnicity								
White Non-Hispanic	541	86.3	(81.3 - 91.3)	34,960	82.6	(81.0 - 84.3)		
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)		
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,215	67.8	(62.9 - 72.8)		
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)		
≥ College Degree	337	87.0	(81.3 - 92.8)	12,236	87.5	(85.0 - 90.1)		
Income								
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)		
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)		
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)		
≥ \$75,000	271	84.6	(77.4 - 91.8)	8,435	88.5	(86.7 - 90.3)		

 $<sup>^{\</sup>rm 1}$  Defined as having had blood cholesterol checked within the past 5 years

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 44a. Cardiovascular disease among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	70,000	7.0	(3.9 - 10.1)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	241	43,400	9.2	(3.4 - 14.9)	14,477	664,600	9.4	(7.5 - 11.4)
Female	402	26,500	5.1	(2.5 - 7.7)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	135	13,700	8.3	(1.8 - 14.9)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	218	38,200	19.8	(14.1 - 25.5)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	550	49,700	7.1	(4.9 - 9.2)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	154	28,900	11.0	(5.4 - 16.6)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	*	*	*	*	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	341	32,300	6.6	(1.2 - 11.9)	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	71	11,700	9.5	(1.2 - 17.8)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	*	*	*	*	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	108	12,500	7.8	(2.9 - 12.7)	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	273	29,800	6.8	(0.7 - 13.0)	8,538	183,500	4.1	(2.9 - 5.3)

 $<sup>^{1}</sup>$  Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 44b. Cardiovascular disease among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	ľ	Nassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	643	6.4	(3.2 - 9.7)	37,810	7.6	(6.4 - 8.7)		
Gender								
Male	241	9.2	(2.9 - 15.6)	14,477	9.7	(7.8 - 11.7)		
Female	402	3.9	(1.9 - 5.9)	23,333	5.6	(4.3 - 6.9)		
Race/Ethnicity								
White Non-Hispanic	550	6.2	(4.3 - 8.2)	35,522	8.2	(7.2 - 9.1)		
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)		
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)		
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)		
Education								
≤ High School	154	9.5	(4.6 - 14.3)	15,486	10.5	(8.0 - 13.0)		
Some College	*	*	*	9,849	6.4	(4.5 - 8.2)		
≥ College Degree	341	6.3	(0.5 - 12.2)	12,393	5.5	(4.2 - 6.9)		
Income								
≤ \$24,999	71	7.7	(1.3 - 14.1)	9,505	12.8	(9.0 - 16.6)		
\$25,000-\$49,999	*	*	*	9,571	6.0	(4.7 - 7.4)		
\$50,000-\$74,999	108	7.1	(2.6 - 11.6)	5,751	8.5	(5.4 - 11.6)		
≥ \$75,000	273	6.8	(0.0 - 13.5)	8,538	4.0	(2.7 - 5.2)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 45a. Coronary heart disease among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	644	60,400	6.1	(3.0 - 9.1)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	241	39,100	8.3	(2.6 - 13.9)	14,479	598,600	8.5	(6.6 - 10.4)
Female	403	21,400	4.1	(1.6 - 6.6)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	*	*	*	*	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	218	31,800	16.5	(11.2 - 21.8)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	551	40,200	5.7	(3.8 - 7.6)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	155	21,500	8.2	(3.1 - 13.2)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	*	*	*	*	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	341	30,600	6.2	(0.8 - 11.6)	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	*	*	*	*	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	*	*	*	*	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	108	9,600	6.0	(2.0 - 10.0)	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	273	28,900	6.6	(0.5 - 12.8)	8,541	151,800	3.4	(2.3 - 4.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 45b. Coronary heart disease among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	ľ	Nassau Co	ounty	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	644	5.7	(2.4 - 8.9)	37,838	6.3	(5.2 - 7.3)		
Gender								
Male	241	8.4	(2.1 - 14.7)	14,479	8.8	(6.9 - 10.8)		
Female	403	3.1	(1.2 - 5.0)	23,359	3.9	(3.0 - 4.8)		
Race/Ethnicity								
White Non-Hispanic	551	5.1	(3.3 - 6.9)	35,552	6.8	(5.9 - 7.7)		
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)		
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)		
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)		
Education								
≤ High School	155	7.1	(2.8 - 11.4)	15,485	8.5	(6.2 - 10.7)		
Some College	*	*	*	9,860	5.5	(3.7 - 7.2)		
≥ College Degree	341	6.1	(0.2 - 12.0)	12,411	4.7	(3.4 - 5.9)		
Income								
≤ \$24,999	*	*	*	9,507	10.7	(7.2 - 14.1)		
\$25,000-\$49,999	*	*	*	9,585	4.7	(3.6 - 5.7)		
\$50,000-\$74,999	108	5.7	(1.8 - 9.6)	5,755	8.0	(4.8 - 11.1)		
≥ \$75,000	273	6.6	(0.0 - 13.4)	8,541	3.3	(2.1 - 4.5)		

 $<sup>^{</sup>m 1}$  Defined as having ever been told by a health professional that respondent had a heart attack or angina

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 46a. Ever diagnosed with diabetes among adults in Nassau County and New York State

		Nassa	au County	<i>'</i>		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	647	67,200	6.7	(4.6 - 8.8)	38,174	1,382,700	9.3	(8.0 - 10.5)		
Gender										
Male	243	18,500	3.9	(1.8 - 6.0)	14,626	530,400	7.5	(6.2 - 8.8)		
Female	404	48,700	9.3	(5.8 - 12.8)	23,548	852,300	11.0	(8.9 - 13.0)		
Age (years)										
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)		
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)		
45-54	*	*	*	*	7,830	279,000	9.8	(6.7 - 12.8)		
55-64	137	21,300	12.7	(5.6 - 19.8)	8,771	332,900	15.5	(12.3 - 18.8)		
≥ 65	219	23,300	12.0	(7.1 - 16.9)	12,645	455,600	17.9	(15.0 - 20.8)		
Race/Ethnicity										
White Non-Hispanic	554	48,800	6.9	(4.7 - 9.0)	35,862	809,300	8.7	(7.8 - 9.5)		
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)		
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)		
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)		
Education										
≤ High School	156	30,700	11.6	(5.7 - 17.4)	15,709	730,000	13.0	(10.5 - 15.6)		
Some College	146	13,300	5.6	(1.7 - 9.5)	9,918	303,500	8.6	(5.9 - 11.2)		
≥ College Degree	342	22,100	4.5	(2.3 - 6.6)	12,464	337,500	6.0	(4.8 - 7.2)		
Income										
≤ \$24,999	*	*	*	*	9,652	543,400	15.5	(11.4 - 19.5)		
\$25,000-\$49,999	*	*	*	*	9,667	291,000	10.0	(7.5 - 12.6)		
\$50,000-\$74,999	109	14,000	8.7	(3.6 - 13.8)	5,777	151,000	7.6	(5.0 - 10.3)		
≥ \$75,000	275	18,700	4.3	(1.9 - 6.6)	8,565	220,800	4.9	(3.8 - 6.1)		

<sup>&</sup>lt;sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant <sup>2</sup> Sample size from survey

 $<sup>^{\</sup>rm 3}$  Estimated number of adults who have diabetes (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 46b. Ever diagnosed with diabetes among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	Nassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	647	5.9	(3.9 - 7.9)	38,174	9.0	(7.8 - 10.3)		
Gender								
Male	243	3.4	(1.5 - 5.2)	14,626	7.5	(6.1 - 8.9)		
Female	404	8.2	(4.8 - 11.6)	23,548	10.4	(8.4 - 12.4)		
Race/Ethnicity								
White Non-Hispanic	554	6.1	(4.0 - 8.2)	35,862	8.4	(7.5 - 9.2)		
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)		
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)		
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)		
Education								
≤ High School	156	10.3	(4.7 - 15.9)	15,709	12.6	(10.0 - 15.2)		
Some College	146	4.7	(1.4 - 8.1)	9,918	8.3	(5.6 - 10.9)		
≥ College Degree	342	4.0	(1.8 - 6.3)	12,464	5.9	(4.7 - 7.2)		
Income								
≤ \$24,999	72	11.6	(1.9 - 21.3)	9,652	15.2	(11.1 - 19.2)		
\$25,000-\$49,999	*	*	*	9,667	9.7	(7.1 - 12.2)		
\$50,000-\$74,999	109	7.4	(3.0 - 11.9)	5,777	7.6	(4.9 - 10.2)		
≥ \$75 <b>,</b> 000	275	4.0	(1.5 - 6.6)	8,565	4.7	(3.6 - 5.9)		

<sup>&</sup>lt;sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when  $\begin{array}{l} \text{pregnant} \\ \text{^{2}} \text{ Prevalence estimates are age-adjusted to the 2000 US Standard population} \end{array}$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 47a. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au County	′		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	649	131,900	13.1	(9.5 - 16.7)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	244	62,900	13.1	(7.2 - 18.9)	14,601	875,800	12.4	(10.1 - 14.7)
Female	405	69,000	13.1	(8.7 - 17.6)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	*	*	*	*	3,931	825,100	18.5	(13.8 - 23.3)
35-44	109	25,900	14.2	(6.1 - 22.4)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	115	18,600	8.6	(3.0 - 14.2)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	137	15,900	9.5	(4.0 - 14.9)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	220	18,200	9.3	(5.1 - 13.6)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	556	95,800	13.4	(9.6 - 17.2)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	156	26,800	10.1	(4.4 - 15.8)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	146	35,200	14.8	(7.4 - 22.3)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	344	69,900	14.0	(8.4 - 19.5)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	*	*	*	*	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	*	*	*	*	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	109	12,200	7.6	(2.1 - 13.2)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	276	51,800	11.7	(7.0 - 16.4)	8,562	577,900	12.8	(10.4 - 15.3)

 $<sup>^{\</sup>mathrm{1}}$  Defined as adults having ever been told by a health professional that they had asthma

 $<sup>^{2}\,\</sup>mathrm{Sample}$  size from survey

 $<sup>^{\</sup>rm 3}$  Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 47b. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	649	14.1	(9.9 - 18.4)	38,096	15.2	(13.3 - 17.0)		
Gender								
Male	244	14.4	(7.8 - 21.1)	14,601	12.6	(10.3 - 14.9)		
Female	405	13.8	(8.4 - 19.2)	23,495	17.5	(14.8 - 20.3)		
Race/Ethnicity								
White Non-Hispanic	556	14.4	(9.9 - 18.9)	35,783	14.9	(13.2 - 16.6)		
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)		
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	156	11.3	(4.2 - 18.4)	15,675	15.1	(12.1 - 18.0)		
Some College	146	16.5	(7.3 - 25.8)	9,893	16.9	(13.4 - 20.4)		
≥ College Degree	344	14.5	(8.3 - 20.7)	12,444	13.3	(10.6 - 16.0)		
Income								
≤ \$24,999	*	*	*	9,643	20.3	(15.0 - 25.6)		
\$25,000-\$49,999	*	*	*	9,641	13.7	(10.2 - 17.2)		
\$50,000-\$74,999	109	7.6	(1.8 - 13.4)	5,762	13.5	(10.0 - 17.0)		
≥ \$75,000	276	12.0	(6.8 - 17.1)	8,562	13.0	(10.6 - 15.4)		

 $<sup>^{\</sup>rm 1}$  Defined as adults having ever been told by a health professional that they had asthma

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 48a. Current asthma among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au County	/		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	645	85,300	8.6	(5.7 - 11.4)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	240	30,000	6.4	(2.6 - 10.1)	14,546	440,700	6.3	(4.8 - 7.7)
Female	405	55,300	10.5	(6.3 - 14.8)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	108	20,100	11.1	(3.6 - 18.6)	4,947	246,800	8.7	(5.8 - 11.6)
45-54	*	*	*	*	7,787	240,000	8.4	(6.1 - 10.8)
55-64	*	*	*	*	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	220	14,100	7.3	(3.3 - 11.2)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	553	66,000	9.3	(5.9 - 12.8)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	156	23,600	8.9	(3.4 - 14.4)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	145	25,200	10.7	(3.7 - 17.6)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	341	36,500	7.4	(3.8 - 11.1)	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	*	*	*	*	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	*	*	*	*	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	273	26,600	6.1	(2.7 - 9.6)	8,541	339,200	7.6	(5.8 - 9.4)

 $<sup>^{\</sup>mathrm{1}}$  Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who currently have asthma (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 48b. Current asthma among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau Co	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	645	9.4	(5.9 - 12.9)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	240	7.2	(2.8 - 11.6)	14,546	6.4	(4.9 - 7.9)
Female	405	11.4	(6.1 - 16.6)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	553	10.4	(6.2 - 14.7)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	156	10.0	(3.1 - 16.9)	15,594	10.8	(8.3 - 13.3)
Some College	145	12.6	(3.6 - 21.7)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	341	7.7	(3.5 - 11.8)	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	*	*	*	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	*	*	*	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	273	6.3	(2.5 - 10.2)	8,541	7.7	(5.9 - 9.5)

 $<sup>^{</sup>m 1}$  Defined as adults having ever been told by a health professional that they had asthma and still have asthma

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 49a. Chronic joint symptoms among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	647	385,900	38.4	(33.6 - 43.2)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	243	156,100	32.5	(25.3 - 39.7)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	404	229,800	43.8	(37.4 - 50.2)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	109	47,400	26.0	(16.4 - 35.7)	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	*	*	*	*	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	*	*	*	*	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	219	104,000	53.6	(46.1 - 61.1)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	554	288,300	40.5	(35.6 - 45.3)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	156	92,500	34.9	(25.2 - 44.6)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	342	183,200	36.7	(30.4 - 43.1)	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	*	*	*	*	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	275	137,000	30.9	(24.6 - 37.2)	8,549	1,874,200	41.7	(38.4 - 45.1)

 $<sup>^1</sup>$  Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 49b. Chronic joint symptoms among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	647	35.1	(30.1 - 40.1)	38,099	42.1	(39.7 - 44.5)		
Gender								
Male	243	29.6	(22.5 - 36.7)	14,595	42.6	(38.8 - 46.3)		
Female	404	40.1	(33.1 - 47.1)	23,504	41.6	(38.6 - 44.7)		
Race/Ethnicity								
White Non-Hispanic	554	37.1	(32.0 - 42.3)	35,785	44.7	(42.7 - 46.6)		
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)		
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,676	43.1	(38.8 - 47.5)		
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)		
≥ College Degree	342	33.1	(26.6 - 39.6)	12,440	40.3	(37.0 - 43.7)		
Income								
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)		
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)		
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)		
≥ \$75,000	275	27.6	(21.4 - 33.7)	8,549	40.8	(37.4 - 44.2)		

 $<sup>^{1}</sup>$  Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 50a. Arthritis among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	645	306,600	30.6	(26.2 - 35.0)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	242	128,300	26.9	(20.3 - 33.6)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	403	178,300	34.0	(28.2 - 39.8)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	109	17,000	9.3	(3.9 - 14.7)	4,972	412,500	14.5	(11.3 - 17.6)
45-54	115	60,800	28.0	(18.6 - 37.4)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	*	*	*	*	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	219	126,200	65.0	(57.7 - 72.3)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	552	243,700	34.4	(29.8 - 39.0)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	*	*	*	*	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	146	83,900	35.3	(25.7 - 44.9)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	340	121,300	24.5	(19.5 - 29.5)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	*	*	*	*	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	273	94,200	21.4	(16.0 - 26.8)	8,536	1,012,800	22.6	(19.8 - 25.3)

<sup>&</sup>lt;sup>1</sup> Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)
<sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 50b. Arthritis among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	645	27.2	(22.7 - 31.6)	38,013	26.6	(24.8 - 28.5)		
Gender								
Male	242	24.6	(18.0 - 31.1)	14,555	23.5	(20.9 - 26.1)		
Female	403	29.5	(23.5 - 35.6)	23,458	29.5	(27.0 - 32.1)		
Race/Ethnicity								
White Non-Hispanic	552	30.4	(25.8 - 35.0)	35,710	30.4	(28.7 - 32.0)		
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)		
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)		
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)		
Education								
≤ High School	*	*	*	15,652	30.4	(26.9 - 33.8)		
Some College	146	31.5	(21.8 - 41.2)	9,874	24.1	(21.0 - 27.1)		
≥ College Degree	340	21.2	(16.5 - 25.9)	12,407	24.7	(21.9 - 27.5)		
Income								
≤ \$24,999	*	*	*	9,618	31.2	(26.3 - 36.1)		
\$25,000-\$49,999	*	*	*	9,624	27.3	(23.2 - 31.3)		
\$50,000-\$74,999	109	32.1	(22.3 - 42.0)	5,753	26.3	(21.9 - 30.7)		
≥ \$75,000	273	18.4	(13.4 - 23.4)	8,536	21.6	(18.8 - 24.3)		

Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)
 Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 51a. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	y		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	180,500	18.1	(14.4 - 21.8)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	242	74,700	15.7	(10.1 - 21.3)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	404	105,900	20.2	(15.4 - 25.0)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	109	22,400	12.3	(5.5 - 19.1)	4,973	418,600	14.5	(11.0 - 18.1)
45-54	115	29,200	13.5	(6.9 - 20.0)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	136	38,900	23.4	(13.8 - 32.9)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	219	63,300	32.8	(25.8 - 39.8)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	554	139,600	19.7	(15.9 - 23.5)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	156	52,800	19.9	(11.5 - 28.4)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	145	48,500	20.6	(12.3 - 28.9)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	342	79,200	16.0	(11.7 - 20.3)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	*	*	*	*	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	*	*	*	*	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	109	41,000	25.6	(15.7 - 35.4)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	275	53,300	12.1	(8.0 - 16.2)	8,544	583,300	13.0	(11.1 - 14.9)

 $<sup>^{\</sup>rm l}$  Defined as activity limitations because of physical, mental or emotional problems

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 51b. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	ľ	Nassau C	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	646	16.9	(13.1 - 20.8)	37,998	20.1	(18.2 - 22.0)		
Gender								
Male	242	15.2	(9.3 - 21.1)	14,566	18.6	(15.8 - 21.4)		
Female	404	18.5	(13.4 - 23.6)	23,432	21.5	(18.9 - 24.1)		
Race/Ethnicity								
White Non-Hispanic	554	18.7	(14.6 - 22.8)	35,688	21.3	(19.7 - 22.9)		
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)		
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)		
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)		
Education								
≤ High School	156	18.8	(9.6 - 28.0)	15,626	23.8	(20.1 - 27.4)		
Some College	145	18.9	(10.2 - 27.7)	9,875	19.3	(16.0 - 22.6)		
≥ College Degree	342	15.2	(10.7 - 19.6)	12,415	16.1	(13.9 - 18.3)		
Income								
≤ \$24,999	*	*	*	9,603	29.7	(24.3 - 35.2)		
\$25,000-\$49,999	*	*	*	9,628	21.2	(16.1 - 26.3)		
\$50,000-\$74,999	109	23.2	(14.0 - 32.4)	5,760	20.1	(15.6 - 24.6)		
≥ \$75,000	275	11.8	(7.6 - 16.0)	8,544	12.7	(10.8 - 14.6)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as activity limitations because of physical, mental or emotional problems

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 52a. Have health problem that requires the use of special equipment among adults in Nassau County and New York State

		Nassa	au Count	y		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	648	65,100	6.5	(4.5 - 8.5)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	243	23,300	4.9	(2.3 - 7.5)	14,617	460,700	6.5	(5.3 - 7.7)
Female	405	41,800	8.0	(4.9 - 11.0)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	*	*	*	*	7,836	182,500	6.4	(4.7 - 8.1)
55-64	137	15,700	9.3	(2.8 - 15.9)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	220	38,000	19.5	(13.4 - 25.7)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	555	51,200	7.2	(4.9 - 9.5)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	156	20,700	7.8	(3.3 - 12.3)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	146	16,200	6.8	(2.9 - 10.8)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	343	28,200	5.7	(3.0 - 8.4)	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	72	10,000	8.1	(0.2 - 15.9)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	89	15,500	11.9	(5.0 - 18.9)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	109	12,100	7.5	(2.4 - 12.7)	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	275	20,000	4.5	(1.9 - 7.2)	8,566	136,000	3.0	(2.1 - 4.0)

 $<sup>^{1}</sup>$  Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone <sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 52b. Have health problem that requires the use of special equipment among adults 1 in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	1	Nassau C	ounty	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	648	5.6	(3.7 - 7.4)	38,171	7.6	(6.6 - 8.6)
Gender						
Male	243	5.1	(2.2 - 8.0)	14,617	6.8	(5.5 - 8.0)
Female	405	6.0	(3.7 - 8.3)	23,554	8.4	(6.9 - 9.9)
Race/Ethnicity						
White Non-Hispanic	555	6.5	(4.2 - 8.8)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
Education						
≤ High School	156	6.6	(2.9 - 10.3)	15,711	10.3	(8.3 - 12.3)
Some College	146	5.4	(2.2 - 8.6)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	343	5.2	(2.4 - 7.9)	12,463	5.3	(4.1 - 6.6)
Income						
≤ \$24,999	72	5.9	(0.3 - 11.4)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	89	9.9	(3.8 - 16.1)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	109	6.6	(2.1 - 11.1)	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	275	4.5	(1.6 - 7.4)	8,566	2.9	(2.0 - 3.9)

 $<sup>^{1}</sup>$  Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 53a. Disability among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	646	193,300	19.3	(15.6 - 23.1)	38,016	3,394,200	22.9	(20.9 - 24.9)		
Gender										
Male	242	78,600	16.5	(10.8 - 22.3)	14,568	1,469,700	20.7	(17.8 - 23.7)		
Female	404	114,700	21.9	(17.0 - 26.9)	23,448	1,924,600	24.8	(22.1 - 27.5)		
Age (years)										
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)		
35-44	109	25,100	13.8	(6.5 - 21.1)	4,974	447,900	15.6	(11.9 - 19.3)		
45-54	115	29,200	13.5	(6.9 - 20.0)	7,811	643,900	22.6	(19.1 - 26.0)		
55-64	136	39,400	23.6	(14.1 - 33.2)	8,740	702,100	33.0	(28.9 - 37.0)		
≥ 65	219	72,900	37.8	(30.5 - 45.1)	12,560	1,010,000	39.9	(36.1 - 43.7)		
Race/Ethnicity										
White Non-Hispanic	554	149,900	21.2	(17.2 - 25.1)	35,703	2,253,400	24.2	(22.6 - 25.9)		
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)		
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)		
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)		
Education										
≤ High School	156	56,200	21.2	(12.7 - 29.7)	15,638	1,544,200	27.6	(23.7 - 31.4)		
Some College	145	53,200	22.6	(14.0 - 31.1)	9,876	755,200	21.4	(17.9 - 24.8)		
≥ College Degree	342	83,800	16.9	(12.5 - 21.4)	12,419	1,021,300	18.2	(15.9 - 20.5)		
Income										
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)		
\$25,000-\$49,999	*	*	*	*	9,629	699,000	24.2	(18.9 - 29.4)		
\$50,000-\$74,999	109	42,500	26.5	(16.6 - 36.4)	5,757	430,100	21.8	(17.2 - 26.4)		
≥ \$75,000	275	57,400	13.0	(8.8 - 17.3)	8,546	632,700	14.1	(12.1 - 16.1)		

 $<sup>^{1}</sup>$  Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

2 Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults with a disability (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 53b. Disability among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	646	18.1	(14.1 - 22.0)	38,016	22.3	(20.4 - 24.3)		
Gender								
Male	242	16.2	(10.1 - 22.2)	14,568	20.9	(18.0 - 23.8)		
Female	404	19.8	(14.6 - 25.1)	23,448	23.6	(21.0 - 26.3)		
Race/Ethnicity								
White Non-Hispanic	554	20.1	(15.8 - 24.3)	35,703	23.6	(21.9 - 25.3)		
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)		
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)		
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)		
Education								
≤ High School	156	19.9	(10.6 - 29.2)	15,638	26.8	(23.0 - 30.6)		
Some College	145	20.5	(11.7 - 29.4)	9,876	20.9	(17.4 - 24.3)		
≥ College Degree	342	16.1	(11.5 - 20.7)	12,419	17.9	(15.5 - 20.3)		
Income								
≤ \$24,999	*	*	*	9,613	33.2	(27.6 - 38.8)		
\$25,000-\$49,999	*	*	*	9,629	23.5	(18.3 - 28.6)		
\$50,000-\$74,999	109	24.0	(14.7 - 33.3)	5,757	21.2	(16.6 - 25.8)		
≥ \$75,000	275	12.8	(8.4 - 17.2)	8,546	13.8	(11.7 - 15.8)		

<sup>&</sup>lt;sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment
<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 54a. Current smoking among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au County	/		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	99,700	10.0	(7.2 - 12.8)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	243	41,600	8.8	(4.5 - 13.0)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	400	58,100	11.1	(7.4 - 14.8)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	108	19,300	10.6	(4.0 - 17.3)	4,977	421,100	14.8	(11.6 - 17.9)
45-54	114	32,500	15.3	(7.7 - 23.0)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	135	16,300	9.8	(3.5 - 16.0)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	218	10,700	5.5	(2.4 - 8.7)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	551	87,700	12.3	(8.8 - 15.9)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	155	31,000	11.7	(5.9 - 17.6)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	145	31,400	13.5	(6.8 - 20.2)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	340	37,400	7.5	(4.0 - 11.0)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	*	*	*	*	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	88	22,000	17.0	(7.9 - 26.2)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	106	21,900	14.3	(5.9 - 22.6)	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	276	42,700	9.6	(5.3 - 13.9)	8,549	468,700	10.4	(8.6 - 12.3)

 $<sup>^{1}</sup>$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^{\</sup>rm 3}$  Estimated number of adult current smokers (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 54b. Current smoking among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau Co	ounty	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	10.1	(7.1 - 13.2)	38,029	17.0	(15.3 - 18.8)
Gender						
Male	243	8.4	(4.2 - 12.7)	14,559	19.2	(16.3 - 22.1)
Female	400	11.7	(7.4 - 16.0)	23,470	15.0	(13.0 - 17.0)
Race/Ethnicity						
White Non-Hispanic	551	12.8	(8.9 - 16.8)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	155	11.2	(5.1 - 17.3)	15,654	22.8	(19.5 - 26.0)
Some College	145	13.8	(6.7 - 20.8)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	340	8.0	(4.0 - 12.0)	12,421	10.0	(7.6 - 12.5)
Income						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	88	17.3	(7.6 - 27.0)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	276	9.1	(4.8 - 13.3)	8,549	10.6	(8.6 - 12.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 55a. Everyday smoking among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au County	1		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	68,600	6.9	(4.4 - 9.4)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	243	28,100	5.9	(2.3 - 9.5)	14,559	848,400	12.0	(10.0 - 14.1)
Female	400	40,600	7.8	(4.4 - 11.2)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	*	*	*	*	4,977	315,200	11.0	(8.4 - 13.7)
45-54	114	29,000	13.7	(6.3 - 21.2)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	*	*	*	*	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	*	*	*	*	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	551	56,600	8.0	(4.9 - 11.0)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	155	23,700	9.0	(3.4 - 14.5)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	145	26,300	11.3	(5.0 - 17.7)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	340	18,600	3.7	(1.1 - 6.4)	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	*	*	*	*	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	88	17,800	13.8	(5.1 - 22.5)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	276	25,900	5.8	(2.2 - 9.5)	8,549	320,200	7.1	(5.5 - 8.8)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

 $<sup>^{2}\,\</sup>mathrm{Sample}$  size from survey

 $<sup>^3</sup>$  Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the

nearest hundred)

4 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 55b. Everyday smoking among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	ľ	Nassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	643	7.0	(4.3 - 9.6)	38,029	11.3	(10.0 - 12.5)		
Gender								
Male	243	5.4	(2.0 - 8.9)	14,559	12.1	(10.1 - 14.1)		
Female	400	8.4	(4.4 - 12.4)	23,470	10.5	(9.0 - 12.0)		
Race/Ethnicity								
White Non-Hispanic	551	8.4	(5.0 - 11.9)	35,729	12.8	(11.5 - 14.1)		
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)		
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)		
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)		
Education								
≤ High School	155	8.8	(3.0 - 14.6)	15,654	16.4	(13.8 - 19.0)		
Some College	145	12.0	(5.2 - 18.7)	9,875	12.1	(9.9 - 14.3)		
≥ College Degree	340	3.9	(0.9 - 6.9)	12,421	5.8	(4.3 - 7.3)		
Income								
≤ \$24,999	*	*	*	9,626	13.4	(10.6 - 16.2)		
\$25,000-\$49,999	88	14.3	(5.0 - 23.5)	9,629	15.3	(12.3 - 18.3)		
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)		
≥ \$75,000	276	5.1	(1.8 - 8.4)	8,549	7.2	(5.5 - 8.9)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 56a. Adults living in homes in which smoking is prohibited<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	646	845,900	84.3	(80.8 - 87.9)	38,045	11,997,100	80.8	(79.0 - 82.6)
Gender								
Male	243	420,000	87.6	(83.1 - 92.1)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	403	425,900	81.4	(76.1 - 86.7)	23,468	6,362,800	82.1	(79.9 - 84.3)
Age (years)								
18-34	68	208,400	85.7	(76.3 - 95.1)	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	109	161,800	89.0	(82.1 - 95.8)	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	115	175,300	80.8	(72.2 - 89.3)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	137	145,200	86.3	(79.7 - 92.9)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	217	155,200	80.6	(75.0 - 86.3)	12,565	1,935,400	76.5	(73.4 - 79.5)
Race/Ethnicity								
White Non-Hispanic	554	577,600	81.1	(76.8 - 85.4)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	154	205,300	77.9	(69.2 - 86.7)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	146	206,500	86.9	(80.4 - 93.3)	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	343	431,500	86.4	(82.0 - 90.9)	12,437	4,817,300	85.7	(83.3 - 88.2)
Income								
≤ \$24,999	71	110,500	89.9	(82.6 - 97.2)	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	*	*	*	*	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	*	*	*	*	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	276	384,000	86.6	(81.0 - 92.2)	8,556	3,883,200	86.5	(84.3 - 88.7)

 $<sup>^{\</sup>rm l}$  Defined as having a rule that prohibits smoking anywhere inside the home

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a \_numerator less than 10)

Table 56b. Adults living in homes in which smoking is prohibited<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	646	84.3	(80.3 - 88.4)	38,045	80.9	(79.2 - 82.7)		
Gender								
Male	243	87.6	(82.8 - 92.4)	14,577	79.4	(76.7 - 82.2)		
Female	403	81.3	(75.1 - 87.6)	23,468	82.3	(80.0 - 84.5)		
Race/Ethnicity								
White Non-Hispanic	554	80.7	(75.7 - 85.7)	35,737	79.9	(78.4 - 81.4)		
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)		
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,647	76.9	(73.8 - 80.0)		
Some College	146	86.7	(80.0 - 93.5)	9,883	79.1	(75.2 - 83.0)		
≥ College Degree	343	87.0	(82.0 - 91.9)	12,437	86.0	(83.5 - 88.5)		
Income								
≤ \$24,999	71	90.4	(82.8 - 97.9)	9,608	74.9	(70.1 - 79.7)		
\$25,000-\$49,999	89	82.5	(72.8 - 92.3)	9,640	79.8	(76.1 - 83.6)		
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)		
≥ \$75,000	276	86.4	(79.8 - 93.0)	8,556	86.6	(84.3 - 88.9)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having a rule that prohibits smoking anywhere inside the home

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 57a. Binge drinking within the past month among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	170,100	17.1	(12.7 - 21.5)	37,745	2,602,000	17.7	(15.7 - 19.8)
Gender								
Male	242	86,100	18.1	(11.3 - 24.8)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	401	84,000	16.2	(10.4 - 21.9)	23,325	889,600	11.6	(9.5 - 13.7)
Age (years)								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	107	20,300	11.4	(3.9 - 18.9)	4,934	584,800	20.6	(16.5 - 24.7)
45-54	114	28,300	13.1	(6.1 - 20.1)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	*	*	*	*	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	*	*	*	*	12,515	88,100	3.5	(2.5 - 4.5)
Race/Ethnicity								
White Non-Hispanic	550	121,800	17.3	(12.8 - 21.8)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
Education								
≤ High School	153	31,600	12.1	(4.4 - 19.8)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	146	48,900	20.6	(10.9 - 30.2)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	342	89,600	18.1	(11.8 - 24.4)	12,358	1,018,700	18.2	(15.3 - 21.1)
Income								
≤ \$24,999	*	*	*	*	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	*	*	*	*	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	276	77,400	17.5	(11.3 - 23.6)	8,501	1,011,400	22.6	(19.7 - 25.6)

 $<sup>^{1}</sup>$  Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 57b. Binge drinking within the past month among adults<sup>1</sup> in Nassau County and New York
State - Age-Adjusted<sup>2</sup>

	r	Nassau C	ounty	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	20.1	(14.9 - 25.3)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	242	19.8	(12.4 - 27.2)	14,420	24.5	(21.0 - 27.9)
Female	401	20.3	(13.1 - 27.6)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	550	20.7	(15.2 - 26.2)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	153	13.1	(4.4 - 21.8)	15,491	17.5	(13.5 - 21.5)
Some College	*	*	*	9,819	18.6	(15.1 - 22.0)
≥ College Degree	342	22.3	(14.7 - 29.8)	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	*	*	*	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	*	*	*	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	276	20.3	(13.0 - 27.6)	8,501	23.2	(20.2 - 26.1)

 $<sup>^{1}</sup>$  Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 58a. Heavy drinking within the past month among adults in Nassau County and New York State

		Nassa	au County	′		New Y	ork State	
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	645	46,000	4.6	(2.7 - 6.5)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	*	*	*	*	14,330	333,300	4.8	(3.9 - 5.7)
Female	403	33,500	6.4	(3.5 - 9.3)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	114	19,100	9.0	(3.2 - 14.8)	7,715	144,600	5.2	(4.1 - 6.2)
55-64	*	*	*	*	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	553	42,300	6.0	(3.5 - 8.4)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	*	*	*	*	15,411	228,100	4.2	(3.2 - 5.2)
Some College	*	*	*	*	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	342	23,300	4.7	(2.2 - 7.2)	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	*	*	*	*	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75 <b>,</b> 000	276	18,300	4.1	(1.7 - 6.5)	8,481	313,900	7.0	(5.4 - 8.6)

 $<sup>^{1}</sup>$  Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month  $^2$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 58b. Heavy drinking within the past month among adults in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau Co	ounty	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	645	4.9	(2.7 - 7.1)	37,553	5.0	(4.3 - 5.8)		
Gender								
Male	*	*	*	14,330	4.9	(3.9 - 5.9)		
Female	403	7.2	(3.6 - 10.8)	23,223	5.1	(4.0 - 6.2)		
Race/Ethnicity								
White Non-Hispanic	553	6.5	(3.6 - 9.4)	35,279	6.4	(5.5 - 7.2)		
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)		
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)		
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)		
Education								
≤ High School	*	*	*	15,411	4.3	(3.2 - 5.3)		
Some College	*	*	*	9,759	4.8	(3.6 - 5.9)		
≥ College Degree	342	5.3	(2.1 - 8.4)	12,306	6.0	(4.5 - 7.4)		
Income								
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)		
\$25,000-\$49,999	*	*	*	9,537	5.8	(4.0 - 7.5)		
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)		
≥ \$75,000	276	4.1	(1.5 - 6.7)	8,481	7.0	(5.4 - 8.7)		

 $<sup>^{1}</sup>$  Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

2 Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	613	673,900	71.2	(66.0 - 76.4)	35,193	8,480,600	61.9	(59.2 - 64.5)
Gender								
Male	231	329,000	72.9	(64.7 - 81.1)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	382	344,900	69.6	(63.1 - 76.1)	21,894	4,123,300	56.8	(53.3 - 60.2)
Age (years)								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	112	175,100	83.1	(75.0 - 91.3)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	128	126,600	79.4	(70.8 - 88.1)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	208	156,900	84.4	(78.2 - 90.5)	11,772	1,908,900	80.3	(76.9 - 83.8)
Race/Ethnicity								
White Non-Hispanic	528	532,100	78.4	(73.7 - 83.1)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	137	172,500	78.9	(70.8 - 86.9)	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	328	343,900	71.9	(64.9 - 78.9)	11,706	3,476,000	65.0	(61.2 - 68.7)
Income								
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	*	*	*	*	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	267	324,300	75.4	(68.4 - 82.4)	8,108	2,934,300	68.2	(64.7 - 71.7)

Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

 $<sup>^{2}\ \</sup>mathrm{Sample}\ \mathrm{size}\ \mathrm{from}\ \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or \_ clinic (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a \_numerator less than 10)

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	Nassau County			N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	613	68.7	(62.9 - 74.6)	35,193	61.4	(58.7 - 64.0)		
Gender								
Male	231	70.6	(61.6 - 79.6)	13,299	67.9	(64.0 - 71.9)		
Female	382	67.0	(59.4 - 74.6)	21,894	55.5	(52.0 - 59.0)		
Race/Ethnicity								
White Non-Hispanic	528	76.4	(70.8 - 82.0)	33,090	67.8	(65.6 - 70.1)		
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)		
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	14,264	59.5	(54.7 - 64.3)		
Some College	137	76.9	(67.8 - 86.0)	9,156	59.7	(54.4 - 65.0)		
≥ College Degree	328	69.6	(61.7 - 77.5)	11,706	64.5	(60.8 - 68.2)		
Income								
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)		
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)		
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)		
≥ \$75,000	267	72.6	(64.4 - 80.8)	8,108	67.5	(64.0 - 71.0)		

Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	372	539,000	89.5	(85.1 - 93.8)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	147	275,900	95.3	(90.1 - 100)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	225	263,100	84.0	(77.4 - 90.7)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	69	99,000	85.8	(76.4 - 95.3)	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	64	107,800	94.4	(89.5 - 99.2)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	67	75,200	93.1	(86.7 - 99.5)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	117	102,500	97.9	(94.7 - 100)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	328	394,300	88.9	(84.3 - 93.4)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	87	149,700	95.9	(90.1 - 100)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	79	113,300	89.8	(80.2 - 99.3)	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	205	275,300	86.2	(79.6 - 92.7)	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	*	*	*	*	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	62	85,700	92.0	(83.8 - 100)	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	174	251,200	89.1	(83.9 - 94.3)	5,771	2,321,600	77.6	(74.0 - 81.2)

<sup>&</sup>lt;sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Nassau County and New York

State - Age-Adjusted<sup>2</sup>

	N	lassau C	ounty	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	372	87.7	(82.5 - 93.0)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	147	95.1	(89.3 - 100)	9,034	81.1	(77.4 - 84.8)
Female	225	81.1	(72.8 - 89.4)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	328	86.5	(80.6 - 92.4)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	87	95.5	(89.2 - 100)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	205	84.1	(76.4 - 91.8)	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	174	87.6	(81.2 - 94.0)	5,771	77.0	(73.4 - 80.6)

<sup>&</sup>lt;sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	628	863,100	88.4	(84.4 - 92.5)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	233	405,700	87.3	(80.9 - 93.6)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	395	457,400	89.5	(84.5 - 94.5)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	*	*	*	*	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	107	168,300	94.5	(89.5 - 99.4)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	113	194,500	91.2	(85.1 - 97.2)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	129	148,200	92.0	(85.4 - 98.6)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	213	149,800	79.7	(73.6 - 85.9)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	538	636,300	91.6	(89.3 - 94.0)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	*	*	*	*	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	146	211,800	89.1	(83.6 - 94.7)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	330	450,400	94.2	(90.0 - 98.5)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	*	*	*	*	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	*	*	*	*	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	107	142,200	89.6	(82.5 - 96.6)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	270	413,800	95.1	(92.5 - 97.7)	8,397	4,184,300	94.4	(92.5 - 96.3)

<sup>&</sup>lt;sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community <sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	lassau C	ounty	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	628	88.7	(84.3 - 93.2)	36,287	89.8	(88.3 - 91.4)
Gender						
Male	233	87.1	(80.2 - 94.0)	13,893	89.1	(86.9 - 91.3)
Female	395	90.2	(84.5 - 96.0)	22,394	90.5	(88.3 - 92.7)
Race/Ethnicity						
White Non-Hispanic	538	92.2	(89.8 - 94.6)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
Education						
≤ High School	*	*	*	14,507	84.6	(81.8 - 87.4)
Some College	146	89.9	(84.7 - 95.2)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	330	94.1	(89.3 - 98.9)	12,142	95.4	(94.0 - 96.7)
Income						
≤ \$24,999	*	*	*	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	*	*	*	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	107	91.5	(85.9 - 97.1)	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	270	95.3	(92.6 - 97.9)	8,397	94.5	(92.6 - 96.4)

 $<sup>^{1}</sup>$  Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 62a. Overweight among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	<b></b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	623	368,200	38.1	(32.7 - 43.4)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	240	251,000	53.4	(45.3 - 61.5)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	383	117,300	23.6	(17.5 - 29.7)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	*	*	*	*	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	*	*	*	*	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	211	67,900	36.4	(29.1 - 43.7)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	531	241,000	35.4	(30.3 - 40.4)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	*	*	*	*	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	329	200,500	41.4	(33.9 - 48.8)	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	*	*	*	*	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	*	*	*	*	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	268	179,800	41.4	(33.5 - 49.4)	8,337	1,659,100	37.5	(34.1 - 40.9)

<sup>&</sup>lt;sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who are overweight (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 62b. Overweight among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	623	37.1	(31.3 - 43.0)	36,710	36.2	(33.8 - 38.6)		
Gender								
Male	240	53.3	(44.7 - 61.8)	14,502	43.9	(40.1 - 47.6)		
Female	383	21.8	(15.1 - 28.4)	22,208	28.9	(25.9 - 31.8)		
Race/Ethnicity								
White Non-Hispanic	531	34.0	(28.6 - 39.4)	34,501	36.0	(34.0 - 38.0)		
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)		
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,079	35.2	(31.1 - 39.4)		
Some College	*	*	*	9,515	38.3	(33.3 - 43.2)		
≥ College Degree	329	40.9	(32.8 - 49.0)	12,062	36.2	(32.7 - 39.8)		
Income								
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)		
\$25,000-\$49,999	*	*	*	9,368	40.6	(34.8 - 46.4)		
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)		
≥ \$75,000	268	40.2	(31.7 - 48.7)	8,337	37.2	(33.7 - 40.6)		

 $<sup>^{1}</sup>$  Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.  $^{2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 63a. Obesity among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	623	168,500	17.4	(13.9 - 21.0)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	240	88,000	18.7	(13.0 - 24.4)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	383	80,500	16.2	(11.9 - 20.4)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	106	29,600	16.5	(8.4 - 24.6)	4,785	662,200	23.6	(19.4 - 27.9)
45-54	111	41,500	20.1	(12.3 - 27.9)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	130	42,900	26.6	(16.6 - 36.5)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	211	35,600	19.1	(12.8 - 25.3)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	531	123,500	18.1	(14.5 - 21.8)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	153	37,900	14.4	(8.0 - 20.8)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	138	47,100	21.7	(13.1 - 30.3)	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	329	82,500	17.0	(12.3 - 21.7)	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	70	15,500	12.8	(3.6 - 22.0)	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	87	19,200	15.8	(5.9 - 25.8)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	268	72,500	16.7	(11.8 - 21.6)	8,337	867,700	19.6	(17.0 - 22.3)

 $<sup>^{1}</sup>$  Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters. <sup>2</sup> Sample size from survey

 $<sup>^{\</sup>rm 3}$  Estimated number of adults who are obese (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 63b. Obesity among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	623	16.1	(12.6 - 19.6)	36,710	23.1	(21.0 - 25.1)		
Gender								
Male	240	16.9	(11.5 - 22.2)	14,502	21.5	(19.0 - 24.0)		
Female	383	15.3	(10.9 - 19.8)	22,208	24.6	(21.5 - 27.8)		
Race/Ethnicity								
White Non-Hispanic	531	17.6	(13.7 - 21.4)	34,501	22.8	(21.0 - 24.5)		
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)		
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)		
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)		
Education								
≤ High School	153	12.5	(6.6 - 18.5)	15,079	29.4	(25.4 - 33.3)		
Some College	138	19.6	(11.6 - 27.6)	9,515	23.1	(19.2 - 27.1)		
≥ College Degree	329	16.3	(11.4 - 21.2)	12,062	16.4	(14.2 - 18.5)		
Income								
≤ \$24,999	70	11.6	(2.5 - 20.7)	9,346	28.3	(22.6 - 34.1)		
\$25,000-\$49,999	*	*	*	9,368	21.0	(17.2 - 24.8)		
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)		
≥ \$75,000	268	15.7	(10.7 - 20.7)	8,337	19.3	(16.7 - 21.9)		

<sup>&</sup>lt;sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 64a. Overweight or obesity among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	623	536,700	55.5	(50.2 - 60.7)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	240	339,000	72.1	(65.0 - 79.2)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	383	197,800	39.7	(33.1 - 46.4)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	*	*	*	*	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	130	107,200	66.4	(57.0 - 75.8)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	211	103,500	55.5	(47.9 - 63.0)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	531	364,500	53.5	(48.3 - 58.7)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	*	*	*	*	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	329	283,000	58.4	(51.5 - 65.3)	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	*	*	*	*	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	*	*	*	*	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	268	252,400	58.1	(50.5 - 65.8)	8,337	2,526,700	57.1	(53.6 - 60.6)

 $<sup>^{1}</sup>$  Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 64b. Overweight or obesity among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	lassau C	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	623	53.2	(47.4 - 59.0)	36,710	59.3	(56.8 - 61.9)		
Gender								
Male	240	70.1	(62.4 - 77.9)	14,502	65.4	(61.3 - 69.4)		
Female	383	37.1	(29.8 - 44.4)	22,208	53.5	(50.1 - 56.8)		
Race/Ethnicity								
White Non-Hispanic	531	51.6	(45.8 - 57.4)	34,501	58.8	(56.7 - 60.8)		
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)		
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,079	64.6	(59.9 - 69.3)		
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)		
≥ College Degree	329	57.2	(49.5 - 64.9)	12,062	52.6	(48.9 - 56.3)		
Income								
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)		
\$25,000-\$49,999	*	*	*	9,368	61.6	(55.1 - 68.1)		
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)		
≥ \$75,000	268	55.9	(47.4 - 64.3)	8,337	56.5	(52.9 - 60.0)		

<sup>&</sup>lt;sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters. <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 65a. Received advice about weight by a health professional among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	623	274,800	28.5	(23.4 - 33.6)	36,729	3,929,700	27.4	(25.1 - 29.7)		
Gender										
Male	240	139,200	29.6	(21.9 - 37.4)	14,499	1,654,800	23.6	(20.6 - 26.6)		
Female	383	135,600	27.4	(20.8 - 34.0)	22,230	2,274,900	31.0	(27.7 - 34.3)		
Age (years)										
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)		
35-44	*	*	*	*	4,779	722,700	25.8	(20.9 - 30.6)		
45-54	110	54,300	26.7	(17.1 - 36.3)	7,490	910,900	33.5	(29.0 - 38.1)		
55-64	*	*	*	*	8,351	741,100	36.1	(32.0 - 40.3)		
≥ 65	212	48,700	26.0	(19.5 - 32.5)	12,313	659,500	26.9	(23.0 - 30.7)		
Race/Ethnicity										
White Non-Hispanic	532	159,800	23.4	(19.5 - 27.4)	34,502	2,386,200	26.5	(24.6 - 28.4)		
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)		
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	15,116	1,469,700	27.5	(23.6 - 31.4)		
Some College	139	59,000	27.2	(17.9 - 36.4)	9,515	946,800	27.8	(23.2 - 32.4)		
≥ College Degree	328	127,900	26.6	(19.8 - 33.3)	12,041	1,445,500	26.2	(23.0 - 29.4)		
Income										
≤ \$24,999	*	*	*	*	9,360	1,079,400	31.4	(25.3 - 37.5)		
\$25,000-\$49,999	*	*	*	*	9,365	696,900	24.6	(19.8 - 29.4)		
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)		
≥ \$75,000	267	100,100	23.2	(16.2 - 30.3)	8,334	1,207,100	27.3	(24.2 - 30.5)		

 $<sup>^{</sup>m 1}$  Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a \_\_numerator less than 10)

Table 65b. Received advice about weight by a health professional among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	N	Nassau C	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	623	26.8	(21.3 - 32.3)	36,729	27.1	(24.8 - 29.4)
Gender						
Male	240	27.7	(19.6 - 35.9)	14,499	23.4	(20.5 - 26.3)
Female	383	25.8	(18.4 - 33.2)	22,230	30.6	(27.3 - 34.0)
Race/Ethnicity						
White Non-Hispanic	532	21.1	(17.2 - 25.0)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,116	27.3	(23.4 - 31.2)
Some College	139	25.9	(15.9 - 35.9)	9,515	27.4	(22.9 - 32.0)
≥ College Degree	328	24.4	(17.2 - 31.5)	12,041	25.9	(22.7 - 29.0)
Income						
≤ \$24,999	*	*	*	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	*	*	*	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	267	20.8	(13.5 - 28.2)	8,334	26.9	(23.7 - 30.0)

<sup>&</sup>lt;sup>1</sup> Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	y	New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	
Total	179	242,300	88.2	(81.9 - 94.5)	9,724	3,412,900	86.8	(83.9 - 89.8)	
Gender									
Male	78	128,900	92.6	(87.0 - 98.1)	3,807	1,468,700	88.8	(85.7 - 91.8)	
Female	*	*	*	*	5,917	1,944,200	85.5	(80.8 - 90.1)	
Age (years)									
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)	
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)	
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)	
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)	
≥ 65	*	*	*	*	2,893	552,300	83.7	(76.9 - 90.6)	
Race/Ethnicity									
White Non-Hispanic	148	142,900	89.4	(84.0 - 94.8)	9,082	2,076,700	87.0	(84.5 - 89.6)	
Black Non-Hispanic	*	*	*	*	*	*	*	*	
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)	
Other Non-Hispanic	*	*	*	*	*	*	*	*	
Education									
≤ High School	*	*	*	*	3,962	1,239,700	84.3	(78.6 - 90.1)	
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)	
≥ College Degree	92	115,000	89.9	(83.1 - 96.6)	3,115	1,290,900	89.3	(85.3 - 93.3)	
Income									
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)	
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)	
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)	
≥ \$75,000	70	91,300	91.2	(84.0 - 98.3)	2,270	1,073,200	88.9	(83.7 - 94.1)	

 $<sup>^{\</sup>mathrm{1}}$  Defined as provided advice by a health professional about their weight and advised to lose weight

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)
<sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	P	Nassau C	ounty	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	179	89.2	(83.5 - 94.9)	9,724	86.9	(83.8 - 89.9)		
Gender								
Male	78	92.5	(87.0 - 98.0)	3,807	88.7	(85.7 - 91.8)		
Female	101	85.8	(75.9 - 95.7)	5,917	85.5	(80.7 - 90.3)		
Race/Ethnicity								
White Non-Hispanic	148	89.2	(83.6 - 94.7)	9,082	87.0	(84.4 - 89.6)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	3,962	84.6	(78.7 - 90.5)		
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)		
≥ College Degree	92	89.8	(82.6 - 96.9)	3,115	89.4	(85.4 - 93.3)		
Income								
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)		
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)		
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)		
≥ \$75,000	70	91.3	(84.1 - 98.4)	2,270	88.7	(83.1 - 94.2)		

<sup>&</sup>lt;sup>1</sup> Defined as provided advice by a health professional about their weight and advised to lose weight

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 67a. No leisure-time physical activity among adults<sup>1</sup> in Nassau County and New York State

		Nassa	au Count	y		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	648	228,400	22.7	(18.1 - 27.4)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	244	91,100	19.0	(12.5 - 25.5)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	404	137,300	26.2	(19.6 - 32.8)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	109	26,800	14.7	(6.6 - 22.9)	4,988	556,000	19.3	(15.0 - 23.6)
45-54	114	53,700	24.8	(15.1 - 34.5)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	137	32,900	19.5	(10.7 - 28.3)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	220	55,600	28.6	(21.7 - 35.5)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	555	150,800	21.2	(16.9 - 25.5)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	*	*	*	*	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	343	68,200	13.7	(9.4 - 17.9)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	*	*	*	*	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	*	*	*	*	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	276	82,200	18.6	(12.6 - 24.5)	8,565	560,800	12.5	(10.3 - 14.6)

Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 67b. No leisure-time physical activity among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	ľ	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	648	22.3	(17.1 - 27.5)	38,128	23.7	(21.5 - 25.9)		
Gender								
Male	244	18.1	(11.5 - 24.7)	14,602	20.9	(17.8 - 23.9)		
Female	404	26.2	(18.5 - 33.9)	23,526	26.3	(23.1 - 29.5)		
Race/Ethnicity								
White Non-Hispanic	555	20.7	(15.9 - 25.5)	35,816	20.0	(18.3 - 21.7)		
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)		
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,683	32.3	(28.3 - 36.4)		
Some College	*	*	*	9,906	22.8	(18.5 - 27.2)		
≥ College Degree	343	13.5	(8.8 - 18.2)	12,458	14.9	(11.8 - 17.9)		
Income								
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)		
\$25,000-\$49,999	*	*	*	9,658	22.0	(18.0 - 26.0)		
\$50,000-\$74,999	*	*	*	5,773	21.6	(16.3 - 26.8)		
≥ \$75,000	276	18.9	(12.2 - 25.7)	8,565	12.6	(10.3 - 14.9)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Nassau County and New York State

	Nassau County					New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	649	282,800	28.1	(23.2 - 33.1)	38,203	4,023,400	27.0	(24.9 - 29.2)		
Gender										
Male	244	113,000	23.5	(15.4 - 31.7)	14,634	1,603,900	22.6	(19.4 - 25.8)		
Female	405	169,700	32.3	(26.3 - 38.4)	23,569	2,419,500	31.1	(28.2 - 34.0)		
Age (years)										
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)		
35-44	*	*	*	*	4,989	688,000	23.9	(19.5 - 28.3)		
45-54	*	*	*	*	7,837	746,800	26.2	(22.3 - 30.1)		
55-64	137	38,600	22.9	(15.3 - 30.5)	8,777	579,100	27.0	(23.6 - 30.5)		
≥ 65	220	61,400	31.5	(24.6 - 38.5)	12,660	747,800	29.4	(26.0 - 32.8)		
Race/Ethnicity										
White Non-Hispanic	556	189,600	26.6	(21.9 - 31.2)	35,884	2,538,000	27.2	(25.4 - 28.9)		
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)		
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)		
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)		
Education										
≤ High School	156	50,700	19.1	(10.7 - 27.6)	15,726	1,309,800	23.4	(19.3 - 27.4)		
Some College	146	77,900	32.8	(22.8 - 42.7)	9,924	964,300	27.2	(23.0 - 31.3)		
≥ College Degree	344	154,200	30.9	(23.5 - 38.2)	12,471	1,747,500	31.0	(27.9 - 34.1)		
Income										
≤ \$24,999	*	*	*	*	9,667	831,500	23.7	(18.6 - 28.7)		
\$25,000-\$49,999	*	*	*	*	9,669	917,200	31.6	(25.6 - 37.6)		
\$50,000-\$74,999	*	*	*	*	5,777	501,000	25.3	(21.4 - 29.3)		
≥ \$75,000	276	130,100	29.3	(21.5 - 37.2)	8,569	1,260,100	28.0	(25.0 - 31.0)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having consumed 5 or more servings of fruits and vegetables in an average day

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	649	28.3	(22.8 - 33.8)	38,203	27.1	(24.9 - 29.2)		
Gender								
Male	244	24.7	(15.9 - 33.4)	14,634	22.7	(19.6 - 25.9)		
Female	405	31.7	(24.6 - 38.7)	23,569	31.0	(28.1 - 34.0)		
Race/Ethnicity								
White Non-Hispanic	556	26.6	(21.2 - 31.9)	35,884	27.1	(25.4 - 28.9)		
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)		
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)		
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)		
Education								
≤ High School	156	19.6	(9.6 - 29.6)	15,726	23.3	(19.3 - 27.3)		
Some College	*	*	*	9,924	27.1	(22.9 - 31.2)		
≥ College Degree	344	31.0	(22.9 - 39.2)	12,471	31.2	(28.0 - 34.3)		
Income								
≤ \$24,999	*	*	*	9,667	23.9	(18.8 - 28.9)		
\$25,000-\$49,999	*	*	*	9,669	32.0	(26.0 - 38.0)		
\$50,000-\$74,999	*	*	*	5,777	25.0	(21.0 - 28.9)		
≥ \$75,000	276	29.7	(21.0 - 38.4)	8,569	28.0	(24.9 - 31.1)		

 $<sup>^{\</sup>rm 1}$  Defined as having consumed 5 or more servings of fruits and vegetables in an average day

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 69a. Had a fall within the past 3 months among adults in Nassau County and New York

		Nassa	au County	У		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	464	62,400	10.9	(7.9 - 13.8)	28,783	1,244,800	17.0	(14.9 - 19.0)		
Gender										
Male	178	27,700	10.4	(6.0 - 14.7)	10,966	489,400	14.4	(11.9 - 16.8)		
Female	286	34,700	11.3	(7.2 - 15.3)	17,817	755,300	19.2	(16.1 - 22.3)		
Age (years)										
18-34	*	*	*	*	*	*	*	*		
35-44	*	*	*	*	*	*	*	*		
45-54	115	14,200	6.6	(2.3 - 10.8)	7,741	445,500	16.0	(12.0 - 20.0)		
55-64	131	19,700	12.0	(6.0 - 18.0)	8,454	336,000	16.5	(13.4 - 19.6)		
≥ 65	218	28,400	14.7	(9.6 - 19.9)	12,588	463,300	18.4	(15.2 - 21.6)		
Race/Ethnicity										
White Non-Hispanic	416	52,100	11.4	(8.0 - 14.7)	27,456	869,000	16.6	(15.1 - 18.1)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)		
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)		
Education										
≤ High School	119	16,500	10.9	(5.5 - 16.2)	12,417	493,000	17.3	(13.3 - 21.3)		
Some College	*	*	*	*	7,232	299,400	18.7	(14.6 - 22.7)		
≥ College Degree	238	36,600	13.3	(8.5 - 18.1)	9,086	441,500	15.4	(12.9 - 17.9)		
Income										
≤ \$24,999	*	*	*	*	7,824	423,900	25.0	(18.5 - 31.6)		
\$25,000-\$49,999	*	*	*	*	7,445	223,600	16.0	(12.8 - 19.2)		
\$50,000-\$74,999	83	15,000	13.8	(6.6 - 21.0)	4,141	151,200	14.6	(10.8 - 18.4)		
≥ \$75,000	188	23,300	9.1	(4.8 - 13.4)	5,874	287,100	12.5	(10.0 - 15.1)		

 $<sup>^{1}</sup>$  Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 69b. Had a fall within the past 3 months among adults<sup>1</sup> in Nassau County and New York
State - Age-Adjusted<sup>2</sup>

	Nassau County			N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	464	10.9	(7.9 - 13.9)	28,783	17.3	(15.0 - 19.5)		
Gender								
Male	178	10.7	(6.3 - 15.0)	10,966	15.0	(12.1 - 17.8)		
Female	286	11.2	(7.1 - 15.3)	17,817	19.4	(16.0 - 22.8)		
Race/Ethnicity								
White Non-Hispanic	416	11.5	(8.1 - 14.9)	27,456	16.9	(15.2 - 18.7)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)		
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)		
Education								
≤ High School	119	11.2	(5.6 - 16.8)	12,417	17.4	(13.0 - 21.7)		
Some College	*	*	*	7,232	19.0	(14.7 - 23.2)		
≥ College Degree	238	13.2	(8.5 - 18.0)	9,086	16.0	(13.1 - 18.8)		
Income								
≤ \$24,999	*	*	*	7,824	25.3	(18.3 - 32.4)		
\$25,000-\$49,999	*	*	*	7,445	15.9	(12.6 - 19.2)		
\$50,000-\$74,999	83	14.3	(6.9 - 21.7)	4,141	14.7	(10.6 - 18.8)		
≥ \$75,000	188	9.2	(4.8 - 13.6)	5,874	12.9	(9.9 - 15.8)		

<sup>&</sup>lt;sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 70a. Had a fall that caused injury within the past 3 months among adults 1 in Nassau County and New York State

		Nassa	au County	•		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	464	19,000	3.3	(1.7 - 4.9)	28,760	335,700	4.6	(3.7 - 5.5)		
Gender										
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)		
Female	286	11,400	3.7	(1.5 - 6.0)	17,803	215,600	5.5	(4.1 - 6.9)		
Age (years)										
18-34	*	*	*	*	*	*	*	*		
35-44	*	*	*	*	*	*	*	*		
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)		
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)		
≥ 65	218	8,400	4.3	(1.5 - 7.2)	12,574	130,300	5.2	(3.4 - 6.9)		
Race/Ethnicity										
White Non-Hispanic	416	14,900	3.3	(1.6 - 4.9)	27,433	270,800	5.2	(4.2 - 6.1)		
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)		
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)		
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)		
Education										
≤ High School	*	*	*	*	12,404	148,900	5.2	(3.6 - 6.8)		
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)		
≥ College Degree	238	9,000	3.3	(1.1 - 5.4)	9,081	101,300	3.5	(2.4 - 4.6)		
Income										
≤ \$24,999	*	*	*	*	7,812	90,200	5.3	(3.6 - 7.1)		
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)		
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)		
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)		

 $<sup>^{1}</sup>$  Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months  $^2$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 70b. Had a fall that caused injury within the past 3 months among adults 1 in Nassau County and New York State - Age-Adjusted<sup>2</sup>

	r	Nassau Co	ounty	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	464	3.3	(1.7 - 4.8)	28,760	4.6	(3.6 - 5.5)		
Gender								
Male	*	*	*	10,957	3.6	(2.4 - 4.7)		
Female	286	3.6	(1.4 - 5.8)	17,803	5.5	(4.1 - 6.9)		
Race/Ethnicity								
White Non-Hispanic	416	3.2	(1.5 - 4.9)	27,433	5.2	(4.2 - 6.2)		
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)		
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)		
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)		
Education								
≤ High School	*	*	*	12,404	5.2	(3.5 - 6.8)		
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)		
≥ College Degree	238	3.3	(1.1 - 5.5)	9,081	3.6	(2.4 - 4.7)		
Income								
≤ \$24,999	*	*	*	7,812	5.1	(3.3 - 7.0)		
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)		
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)		
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)		

<sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)